



Level of Awareness and Knowledge of Folic Acid Intake among Women in the Reproductive Age in ARAR Region, Saudi Arabia

**Abdulrahman Mazki Alanazi^{a*}, Abdullah Juayf Alanazi^{a#},
Yasir Wadi Alanazi^{a#}, Muhammad Sari Alanazi^{a#}, Ghada Rezk^{b#}
and Sabry Mohamed Hammad^{c,d#}**

^a Saudi Board of Family Medicine, Joint Program, Northern Borders Region, Kingdom of Saudi Arabia.

^b Egyptian Board, Family Medicine Consultant, Northern Border, Salhya Awasad Primary Health Care, KSA.

^c Faculty of Medicine, Mansoura University, Egypt.

^d Northern Border, General Health Affairs, KSA.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i58A34118

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/79482>

Original Research Article

**Received 06 October 2021
Accepted 12 December 2021
Published 15 December 2021**

ABSTRACT

Background: Folic acid reduces the risk of neural tube defects. As approximately 50% of pregnancies are unintended, women of reproductive age should be aware of the importance of folic acid. This study aims to assess the level of awareness and knowledge among childbearing female on the importance of preconception folic acid supplementation in preventing neural tube defects (NTDs) in Arar city.

Methodology: A population-based cross-sectional study was conducted in Arar city, Northern Saudi Arabia on included Saudi women in reproductive age. A pre-designed online questionnaire was used for data collection. Data was entered and analyzed using statistical package for Social Sciences Program, version 20 (SPSS Inc., Chicago, Illinois, USA). The results were displayed as

[#] Resident Family Medicine Physician;

^o Consultant of Family Medicine;

[#] Professor of Public Health & Community Medicine;

*Corresponding author: E-mail: r7moni_330_@hotmail.com;

counts, percentages. Appropriate statistical tests will be used for comparisons. $P \leq 0.05$ is considered significant.

Results: The sample consisted of 428 women in reproductive age, 60.1% aged between 30 to 40 years, 39% aged between 20 to 30 years and only 0.9% aged between 18 to 20 years old. 91.8% were married. 85% of participants reported that their doctor prescribed folic pills for them before. 95.1% think that women should take folic acid pills in pregnancy while 75.2% think that it is necessary to take folic pills pre-pregnancy. 85.3% had taken folic acid pills in previous pregnancy. 31.3% think that it should be stopped after pregnancy. 84.6% think that folate deficiency causes fetal abnormalities. Only 8.4% of studied women reported attending seminar or lecture on folic acid importance.

Conclusion: The results of the current study affirmed good knowledge of folic acid supplementation usage during pregnancy among women in Arar, KSA. There was a significant association between knowledge of folic acid taking importance during pregnancy with marital status, number of children, and educational level of participants.

Keywords: Folic Acid; women; reproductive age; pregnancy; Saudi Arabia.

1. INTRODUCTION

Neural tube defects (NTDs), including spina bifida, anencephaly, encephalocele are most common reasons of morbidity and mortality among infants and neonates. The incidence of NTDs has decreased over the past 60 years and is about 1-2 per 1000 babies in the general population [1].

There is strong evidence that folic acid intake by women in reproductive age improves normal embryonic development and prevents NTDs [2]. Although, the exact mechanism of NTD prevention by folic acid remains largely unknown, some studies show evidence of increase in NTD during exposure to folate antagonist medications and the presence of high anti-bodies titre to folate receptors in women with a past history of NTD [3,4].

Although, folic acid plays a role in preventing anemia and peripheral neuropathy in mother during pregnancy, it can also prevent growth retardation, congenital heart defects, low birth weight, smaller chest and head circumference, cleft lip and preterm delivery of the fetus [5,6].

Normal diet is very important for pregnant women. It contains proteins, fats, carbohydrates, vitamins, and minerals. It is the best way to receive nutrients, but basic vitamin administration can also have a beneficial role, in case of unsuspected deficiency. vitamin supplementation does not replace a balanced diet but rather confirm that a pregnant woman is taking enough daily requirements [7].

Folic Acid can be found in a lot of foods such as eggs, meat, liver, and leafy vegetable. These source alone may not contain the ideal recommended daily of 400 mcg, since some of the folate sources can be lowered during cooking it [8,9,10,11].

A study conducted in Texas, USA, showed 78% of women had good awareness towards folic acid supplementation, 28% of participants believed it prevented birth defects and 25% knew to take it before conception. Only 33% of respondents reported daily folic acid supplementation [12].

Similar study in Lebanon showed that only 60% of women had heard of folic acid and only 14% of participants had good knowledge regarding the benefits of folic acid to prevent NTDs [13]. Another study in Qatar displayed that 53.7% of studied subjects had heard of folate and only 20.3% of respondents reported their actual intake of folic acid [14].

Regarding the level of awareness toward preconception folic acid supplementation among female college students, a study was conducted in Jeddah, KSA, that showed only 12% knew the benefit of folic acid in preventing NTDs. A previous research also displayed that the majority of women in the Saudi Arabia have low level of knowledge scores for good nutrition during pregnancy and lactation [15,16].

In addition, a study done in Hail Region, Saudi Arabia showed high awareness among women, toward folic acid supplementation, as 91% had heard or read about FA and 81% knew that FA has an important role in preventing NTDs, moreover 70% had good knowledge about

proper timing for FA supplementation and believed that FA should be taken during the first trimester of pregnancy, whereas 41 % responded believed that it should be taken before conception, while only 10% of the participants believed that it should be taken before conception as well as during pregnancy. Over 77% of subjects obtained their information of FA intake from medical care providers including physicians, nurse, dietitian, and midwives [17].

1.1 Study Rationale

Reviewing the literature, showed that there is no study done in Northern Border Region in Kingdom of Saudi Arabia, which explored awareness, knowledge, attitude and behavior regarding Folic Acid Intake among women in the reproductive age in Arar, Saudi Arabia. This study aims to assess prevalence of intake of folate and perform a KAP on folic acid intake among women in reproductive age in Arar Region, Saudi Arabia.

1.2 Aim of the Study

The aim of this study to assess the level of awareness and knowledge among childbearing female on the importance of preconception folic acid supplementation in preventing neural tube defects (NTDs) in Arar city.

2. METHODOLOGY

2.1 Subjects and Methods

Study design: A cross sectional study was conducted in 2021.

Study setting: Arar city, Norther Saudi Arabia.

Study population: Saudi women in reproductive age in Arar city, KSA

2.2 Inclusion Criteria

- Saudi women
- Reproductive age group aged 18 -40 years old
- Live in Arar city
- Welling to participate

2.3 Exclusion Criteria

- Non-Saudi women
- Men
- Younger than 18 years old or older than 40
- Live outside Arar city

- Health care workers
- Don't want to participate

2.4 Sampling

The sample size is calculated based on the sample size formula [7]. $n = [z/ \alpha]^2 \times p [1-p]$, where n is the sample size, z is the confidence interval (CI) assumed to be 1.96, α is assumed to be 0.05 . p is the probability according to a previous study conducted in Saudi Arabia that reported 68% level of awareness regarding folic acid supplementation. The minimum sample size calculated for the present study is 335 women.

2.5 Sampling Technique

Convenient sample from female responding to electronic questionnaire.

2.6 Data Collection Tool

Self-administer questionnaire was distributed online. The questionnaire is developed by researcher after literature review. It contains questions on demographic data (e.g: age ,education ,income).Twelve questions on knowledge (e.g: importance of folic acid supplementation).Three questions on attitude and three questions on practice .The questionnaire was reviewed by three experts for validity and reliability.

2.7 Data Management

Data was entered and analyzed using statistical package for Social Sciences Program, version 20 (SPSS Inc., Chicago, Illinois, USA). The results were displayed as counts, percentages. appropriate statistical tests will be used for comparisons. $P \leq 0.05$ is considered significant.

3. RESULTS

According to table 1; the sample consisted of 428 women in reproductive age, 60.1% aged between 30 to 40 years, 39% aged between 20 to 30 years and only 0.9% aged between 18 to 20 years old. Most study participants 87.6% had university education while only 1.4% were uneducated. 91.8% were married. 50.5% of all participants were working.

Table 1. Sociodemographic characteristics of participants (n=428)

Parameter		No.	Percent
Age	• 18- 20 years old	4	0.9
	• 20 - 30 years old	167	39.0
	• 31 - 40 years old	257	60.1
Education level	• Uneducated	6	1.4
	• Preparatory	9	2.1
	• secondary	38	8.9
	• university	375	87.6
Job	• work	216	50.5
	• not work	212	49.5
Social status	• married	393	91.8
	• divorced	23	5.4
	• widow	12	2.8

Table 2. Number of children, pregnancy and planning for pregnancy, history of miscarriage and having a child with malformation (n= 428)

Parameter		No.	Percent
Number of children, if any	• None	69	16.1
	• 1	56	13.1
	• 2	60	14.0
	• 3	44	10.3
	• 4	50	11.7
	• 5	54	12.6
	• 6	56	13.1
	• More than 7	39	9.1
Current pregnancy	• Yes	63	14.7
	• No	365	85.3
Future plans for pregnancy	• Yes	220	51.4
	• No	208	48.6
History of miscarriage	• Yes	160	37.4
	• No	268	62.6
Child with malformations	• Yes	8	1.9
	• No	420	98.1

In Table 2; 16.1% had no children while 13.1% had one child, 14% had two, 10.3% had three, 11.7% had four, 12.6% had five and 13.1% had six children. 14.7% of all women were pregnant at the time of the study while 51.4% have plans for future pregnancy. 37.4% reported history of miscarriage. Only 1.9% had a child with malformation.

As illustrated in Table 3, 4 85% of participants reported that their doctor prescribed folic pills for them before. 83.2% of women know there is lot of information now available on the use of folic before pregnancy. 90% know that iron deficiency causes anemia. 95.1% think that women should take folic acid pills in pregnancy while 75.2%

think that it is necessary to take folic pills pre-pregnancy. 96.3% think that folic acid pills are important in the first trimester of pregnancy, 2.3% think it is important in the second trimester while 1.4% think it is important in the third trimester.

95.3% reported that doctors recommend taking folic pills. 85.3% had taken folic acid pills in previous pregnancy. 31.3% think that it should be stopped after pregnancy. 15.7% think that folic acid dose is three pills per day. 84.6% think that folate deficiency cause fetal abnormalities.

Only 10.3% think that joint pain is a side effect of folic acid pills. Regarding natural sources of folic, 89.5% reported eggs and 10.5% bread. Only

52.8% reported that their diet is rich with folates. 75% think that folic acid is a vitamin, 16.6% think it is iron and only 8.4% think it is protein. Only 8.4% of studied women reported attending seminar or lecture on folic acid importance.

According to Table 5, there was a significant association between knowledge of folic acid taking importance during pregnancy with marital status, number of children, and educational level of participants.

Table 3. Knowledge of participants of importance of folic acid during pregnancy (n=428)

Parameter		No.	Percent
Previous Folic acid prescription by a doctor	• Yes	364	85.0
	• No	64	15.0
There is a lot of information now available on the use of folic before pregnancy	• Yes	356	83.2
	• No	72	16.8
Women should take folic pills during pregnancy	• Yes	407	95.1
	• No	21	4.9
Folic pills are important during	• The first three months	412	96.3
	• middle three months	10	2.3
	• the last three months	6	1.4
Doctors recommend taking folic pills	• Yes	408	95.3
	• No	20	4.7
Pills should be stopped after pregnancy	• Yes	134	31.3
	• No	294	68.7
Iron deficiency lead to anaemia	• Yes	385	90.0
	• No	43	10.0

Table 4. Knowledge of participants of folic acid and its associated factors and education about it (n=428)

Parameter		No.	Percent
Folic acid dose is three pills per day	• Yes	67	15.7
	• No	361	84.3
Joint pain is one of the side effects of folic acid	• Yes	44	10.3
	• No	384	89.7
Natural sources of folic	• eggs	383	89.5
	• the bread	45	10.5
There is a preferred time during the day to take folic	• Yes	149	34.8
	• No	279	65.2
It is necessary to take folic pills in the pre-pregnancy period	• Yes	322	75.2
	• No	106	24.8
Overcooking reduce the nutritional value of folic acid	• Yes	172	40.2
	• No	256	59.8
Folate deficiency cause foetal abnormalities	• Yes	362	84.6
	• No	66	15.4
Folic pills are	• protein	36	8.4
	• iron	71	16.6
	• vitamins	321	75.0

Parameter		No.	Percent
History of taking folic pills in a previous pregnancy	• Yes	365	85.3
	• No	63	14.7
Attend any seminar or lecture on folic acid importance	• Yes	36	8.4
	• No	392	91.6
Have diet rich in folate	• Yes	226	52.8
	• No	202	47.2

Table 5. Association between participants knowledge of folic acid and sociodemographic characteristics of participants

		Women should take folic acid during pregnancy		Total (N=428)	P value
		Yes	No		
Age	Less than 20	3	1	4	0.278
		0.7%	4.8%	0.9%	
	20 - 30 years old	157	10	167	
		38.6%	47.6%	39.0%	
	31 - 40 years old	166	8	174	
		40.8%	38.1%	40.7%	
41 – 50 years old	76	2	78		
	18.7%	9.5%	18.2%		
Education level	uneducated	5	0	5	0.005
		1.2%	0.0%	1.2%	
	Preparatory	6	0	6	
		1.5%	0.0%	1.4%	
	secondary	7	2	9	
		1.7%	9.5%	2.1%	
university	33	5	38		
	8.1%	23.8%	8.9%		
Working status	work	361	14	375	0.107
		88.7%	66.7%	87.6%	
	not work	209	7	216	
Social status	married	51.4%	33.3%	50.5%	0.001
		198	14	212	
	48.6%	66.7%	49.5%		
	divorced	378	15	393	
		92.9%	71.4%	91.8%	
widow	21	2	23		
Number of children, if any	Nothing	5.2%	9.5%	5.4%	0.037
		8	4	12	
	2.0%	19.0%	2.8%		
	1	60	9	69	
		14.7%	42.9%	16.1%	
	2	55	1	56	
		13.5%	4.8%	13.1%	
	3	59	1	60	
		14.5%	4.8%	14.0%	
4	43	1	44		
	10.6%	4.8%	10.3%		
5	48	2	50		
	11.8%	9.5%	11.7%		
6	51	3	54		
	12.5%	14.3%	12.6%		
	55	1	56		

		Women should take folic acid during pregnancy		Total (N=428)	P value
		Yes	No		
	More than 7	13.5%	4.8%	13.1%	
		36	3	39	
		8.8%	14.3%	9.1%	
Future plans for pregnancy	Yes	210	10	220	0.722
	No	197	11	208	
		51.6%	47.6%	51.4%	
History of previous miscarriages	Yes	156	4	160	0.075
	No	251	17	268	
		48.4%	52.4%	48.6%	
Child with malformations	Yes	8	0	8	0.517
	No	399	21	420	
		61.7%	81.0%	62.6%	
		2.0%	0.0%	1.9%	
		98.0%	100.0%	98.1%	

4. DISCUSSION

In addition to the birth defects, FA deficiency during pregnancy may increase the risk of preterm birth, low birth weight and intrauterine growth retardation of the fetus, and it may raise the level of homocysteine in the blood. High homocysteine levels may lead to spontaneous abortions, placental abruption and pre-eclampsia [5]. This study aims to assess the level of awareness and knowledge among childbearing female on the importance of preconception folic acid supplementation in preventing neural tube defects (NTDs) in Saudi Arabia.

In our study, 95.1% think that women should take folic acid pills in pregnancy and 85.3% had taken folic acid pills in previous pregnancy. Another Saudi study reported that, during the pregnancy, almost 95% of women were using these supplementations to avoid maternal (anemia, peripheral neuropathy) and fetal (congenital abnormalities and NTD) complications [14]. Alreshidi et al.[18] studies have conducted the survey from Riyadh city and confirms 42.2% of pregnant women knows about supplementation of FA. Almost 80% of were known about prevention of NTDs and 46.8% of pregnant women consumed FA during pregnancy. Alodan et al.[19] studies performed at Security forces hospitals in Riyadh city and concluded as pregnant women had 80% of knowledge about FA supplementation but 53% of women only consumed the FA during pregnancy. Al-Rakaf et al [20]. also performed a similar study in pregnant women from Prince Sultan Hospital in Riyadh and confirmed the same conclusion from Alreshidi and Alodan et al [18,19]. studies. An

earlier research by Albader et al.[21] carried out in Riyadh city and AbdRabou et al.[22] studies carried out in Sakaka city were also confirms the similar conclusions from the prior studies carried in Saudi Arabia. Ahmad et al.[23] from Taibah University in Madinah and Al-Holy et al.[24] in the University of Hail in Al-Hail performed a questionnaire-based study and concluded as women had limited knowledge about FA supplementations in married and unmarried women. Maximum studies implemented in different cities in Saudi Arabia has found the similar confirmation as pregnant women have knowledge about FA supplementation and almost 50% of women were not achieving and further effects the NTDs for their neonates [14].

Outside Saudi Arabia, lower rates were reported in previous literature 76.59% reported hearing about FA and 93.89% took it during pregnancy [25]. A cross-sectional survey in Turkey found that 48.2% of the participants were aware that FA is necessary for the prevention of congenital anomalies. Even more significantly, although more than 88.2% of pregnancies were planned among women, only 14.2% of them reported using FA in the preconception period [26,27], Maher et al. showed that nearly 98% of women reported hearing about FA [28]. Lower rates were reported in France as 36% of women never heard of folic acid and 82% were not aware of its benefits [29]. In Turkey, another survey by Köken, Güleğül N et al. reported that (42.2%) of participants heard about folic acid and (48.2%) of them stated that folic acid could prevent birth defects [26].

Folic acid supplementation before pregnancy is known to significantly reduce the risk of having a baby with neural tube defects (NTDs). Therefore, it is important for women to be aware of the effects of folic acid supplementation before pregnancy. Starting FA supplementation before conception reduces the incidence of NTDs by 50–70% [30]. In our study, 75.2% of participants know that it is necessary to take folic pills pre-pregnancy. A previous study reported lower rates as only 10.3% of 1,277 pregnant women reported taking folic acid during the periconceptional period [31]. Comparable results were reported by Köken, Güleğül N et al., only 14.2% of them stated that they had used folic acid in the pre-conception period [26]. Another study reported higher results as 66.41% of participants took it before pregnancy [27]. A high proportion of women worldwide, in both developed and developing countries, needs to be educated that FA should be taken before pregnancy and continued during the first three months after conception [32].

Folic acid supplementation was reported to have a significant positive association with education level, being employed, and planned pregnancy. However, other variables, including age, husband's age, husband's education and employment status, the number of prior pregnancies, economic satisfaction, and household food security, were nonsignificant factors affecting folic acid supplementation [33]. These findings are consistent with studies by Riazi et al. [34], Roth et al. [35], and De Santis et al. [36]. It has been shown that higher education levels in women can increase their awareness about the necessity of folic acid supplementation, and their understanding of the important period of folic acid supplementation [37].

Regarding factors associated with knowledge, our study indicated a significant association between knowledge of folic acid taking importance during pregnancy with marital status, number of children, and educational level of participants. Another study reported that knowledge was higher in older women, women in a couple and women with higher educational level [29].

Even with all these health benefits of folic acid supplementation, recent findings shows that higher population exposure to folic acid may have a negative impact on certain developmental and degenerative disorders. For example, there is currently a lot of focus on the role of folic acid

fortification in increasing the risk of colon cancer. During pregnancy, a similar, possibly unanticipated negative effect has been observed. Synthetic folic acid inhibits human dihydrofolate reductase, resulting in unmetabolized folate in the bloodstream, which may mask irreversible pernicious anaemia caused by B12 deficiency. Low maternal vitamin B12 and high folic acid status during pregnancy may contribute to offspring insulin resistance and obesity [38].

5. CONCLUSION

The results of the current study affirmed good knowledge of folic acid supplementation usage during pregnancy among women in Arar, KSA. There was a significant association between knowledge of folic acid taking importance during pregnancy with marital status, number of children, and educational level of participants. Public health interventions among Saudi women in reproductive age to improve awareness of FA's roles before and during pregnancy are crucial to curtail birth defects.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

CONSENT AND ETHICAL APPROVAL

Informed consent was taken from all study participants and authors assured them that Confidentiality of their data maintained during the study. Research clearance and approval will be obtained from research ethical committee of Northern Borders General Health Affairs.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Laurence KM. The genetics and prevention of neural tube defects and 'uncomplicated'

- hydrocephalus. In: Emery AEH, Rimoin DL, eds. *Principals and practice of medical genetics*, 2nd ed. New York, Churchill Livingstone. 1990:323
2. Finnell RH, Shaw GM, Lammer EJ, Brandl KL, Carmichael SL, and Rosenquist TH. Gene-nutrient interactions: importance of folates and retinoids during early embryogenesis. *Toxicology and Applied Pharmacology*. 2004;198(2):75-85
 3. Vergel, RG, Sanchez LR, Heredero BL, Rodriguez, PL, Maritnez, AJ. Primary prevention of neural tube defects with folic acid supplementation: Cuban experience. *Prenat. Diagn.* 1990;10 (3):149-152.
 4. Brough, L., Rees, G.A., Crawford, M.A., Dorman, E.K.,. Social and ethnic differences in folic acid use preconception and during early pregnancy in the UK: effect on maternal folate status. *J. Hum. Nutr. Diet.* 2009;22 (2):100-107.
 5. Greenberg JA, Bell SJ, Guan Y, Yu YH. Folic Acid supplementation and pregnancy: More than just neural tube defect prevention. *Rev Obstet Gynecol* . 2011 ;4:52-9.
 6. Hassan H. Call For Change Level Of Knowledge, Awareness And Attitude To Follow A High Folate Diet Among Pregnant Women. *Journal of Nursing and Health Science*. 2016;5(1):93-100. www.iosrjournals.org.
 7. The American Congress of Obstetricians and Gynecologists (ACOG). *Essential Nutrients and Vitamins for Pregnancy*; 2011. Available:http://www.americanpregnancy.org/pregnancyhealth/nutrient_svitaminspregnancy.html
 8. Brouwer IA, Van Dusseldorp M, Thomas C M, Duran M, Hautvast J G, Eskes T K, and Steegers-Theunissen R P .. Low-dose folic acid supplementation decreases plasma homocysteine concentrations: a randomized trial. *The American Journal of Clinical Nutrition*. 1999; 69(1): 99-104
 9. Cuskelly GJ, McNulty H, and Scott JM. Fortification with low amounts of folic acid makes a significant difference in folate status in young women: implications for the prevention of neural tube defects. *The American Journal of Clinical Nutrition*. 1999;70(2): 234-239.
 10. Werler M M, Louik C, and Mitchell A A. Achieving a public health recommendation for preventing neural tube defects with folic acid. *American Journal of Public Health*. 1999;89(11): 1637-1640.
 11. Fatima A, Saif N, Ghani M, Ali S. Knowledge of women; supplementation of folic acid in perconceptional period. *Professional Medical Journal*. 2016;1:23(7).
 12. Canfield MA, Przybyla SM, Case AP, Ramadhani T, Suarez L, Dyer J. Folic acid awareness and supplementation among Texas women of childbearing age. *Prev Med (Baltim)*. 2006;43(1):27-30.
 13. Claudine NH, Maya J, Mohamad S, Salim MA Awareness and intake of folic acid for the prevention of neural tube defects among Lebanese women of childbearing age. *Matern Child Health J*. 2012;16:258-265.
 14. Abdulbari 8, Mohammed GA, Al Maadid, Daoud AE, Al-Bast, et al.. Maternal knowledge, attitude and practice on folic acid intake among Arabian Qatari women. *Reproductive Toxicology*. 2006;21:21-25.
 15. Kari JA, Bardisi ES, Baitalmal RM and Ageely GA. Folic Acid Awareness among Female College Students: Neural Tube Defects Prevention. *Saudi Medical Journal*. 2008;29:1749-51.
 16. Mansour A A and Hassan SA. Factors that Influence Women's Nutrition Knowledge in Saudi Arabia," *Health Care for Women International*, 1994;15(3):213-23.
 17. Al-Holy M, Epuru S, Abu-Jamous D, Ashankyty I. Awareness of Folic Acid Intake among Women in the Childbearing Age in Hail Region-Saudi Arabia. *Food Nutr Sci [Internet]*. Available:<http://dx.doi.org/10.4236/fns.2013.41008%0Ahttp://www.scirp.org/journal/fns>
 18. Alreshidi FF, Almujiil AS, Malak AS. Awareness of folic acid use among Saudi women attending outpatient clinics at King Fahad Medical City. *J Family Med Prim Care*. 2018;7:957-62.
 19. AlOdan AA, Ghoraba DA. Maternal knowledge and use of folic acid among Saudi women attending antenatal care clinic at Security Forces Hospital, Riyadh, Saudi Arabia. *IOSR J Nurs Health Sci*. 2018;7(5):11-9.
 20. Al Rakaf MS, Kurdi AM, Ammari AN, Al Hashem AM, Shoukri MM, Garne E, et al. Patterns of folic acid use in pregnant Saudi women and prevalence of neural tube defects—Results from a nested case—

- control study. *Prev Med Rep.* 2015;2: 572–6.
21. Albader NA, Negm DR, El Gabry EK, El-Sayed MM, Arzoo S. Knowledge and practice of folic acid supplementation and impact of income level on awareness among women of child-bearing age in Saudi Arabia. *Trop J Pharm Res.* 2019;18:1323–30.
 22. AbdRabou MAJIJoP. Awareness of folic acid intake for prevention of neural tube defects among women in Sakaka, Saudi Arabia. 2019;15:274–9.
 23. Ahmad B, Anam N, Khalid N, Mohsen R, Zaal L, Jadidy E, et al. Perceptions of women of reproductive age about vitamin and folic acid supplements during pregnancy, Taibah University, Almadinah Almunawwarah, Kingdom of Saudi Arabia. *J Taibah Univ Med Sci.* 2013;8:199–204.
 24. Al-Holy M, Eideh Aa, Epuru S, Abu-Jamous D, Ashanky I. Awareness of folic acid intake among women in the childbearing age in hail region—Saudi Arabia. *Food Nutr Sci.* 2013;4:49–55.
 25. Medawar G, Wehbe T, Jaoude EA. Awareness and Use of Folic Acid among Women of Childbearing Age. *Ann Glob Health.* 2019;85(1):54. Published 2019 Apr 9. DOI:10.5334/aogh.2396
 26. Koken G, Derbent A, Erol O, et al. Awareness and use of folic acid among reproductive age and pregnant women Department of Obstetrics and Gynaecology. Antalya Education and Research Hospital, Antalya, Turkey; 2013.
 27. Ahmad B, Anam N, Khalid N, et al. Perceptions of women of reproductive age about vitamin and folic acid supplements during pregnancy, Taibah University, Almadinah Almunawwarah, Kingdom of Saudi Arabia. *Journal of Taibah University Medical Sciences.* 2013;8(3): 199–204. DOI: 10.1016/j.jtumed.2013.08.002
 28. Maher M and Keriakos R. Women's awareness of periconceptional use of folic acid before and after their antenatal visits. *Clin Med Insights Women's Health.* 2014;1(7):9–15. DOI: 10.4137/CMWH.S13535
 29. Salgues M, Damase-Michel C, Montastruc JL, Lacroix I. Connaissance des femmes en âge de procréer sur l'acide folique : des progrès à faire ! [Women's knowledge of folic acid]. *Therapie.* 2017;72(3): 339–343. Available: <https://doi.org/10.1016/j.therap.2016.10.001>
 30. Kim J, Yon M, Kim CI, et al. Preconceptional use of folic acid and knowledge about folic acid among low-income pregnant women in Korea. *Nutr Res Pract.* 2017;11(3): 240-246. DOI:10.4162/nrp.2017.11.3.240
 31. Kim MH, Han JY, Cho YJ, Ahn HK, Kim JO, Ryu HM, Kim MY, Yang JH, Nava-Ocampo AA. Factors associated with a positive intake of folic acid in the periconceptional period among Korean women. *Public Health Nutr.* 2009;12: 468–471.
 32. Wang M, Li K, Zhao D and Li L. The association between maternal use of folic acid supplements during pregnancy and risk of autism spectrum disorders in children: A meta-analysis. *Mol. Autism.* 2017; 2(8): 51. DOI: 10.1186/s13229-017-0170-8
 33. Ezzeddin N, Zavoshy R, Noroozi M. Prevalence of folic acid supplement consumption before and during pregnancy, and its determinants among community health center referrals. *Obstet Gynecol Sci.* 2019;62(6): 454-461. DOI:10.5468/ogs.2019.62.6.454
 34. Riazi H, Bashirian S, Cheraghi F. Folic acid Intake during preconceptional period. *Avicenna J Nurs Midwifery Care.* 2010;18:38–48.
 35. Roth C, Bjørke-Monsen AL, Reichborn-Kjennerud T, Nilsen RM, Smith GD, Stoltenberg C, et al. Use of folic acid supplements in early pregnancy in relation to maternal plasma levels in week 18 of pregnancy. *Mol Nutr Food Res.* 2013;57:653–660.
 36. De Santis M, Quattrocchi T, Mappa I, Spagnuolo T, Licameli A, Chiaradia G, et al. Folic acid use in planned pregnancy: an Italian survey. *Matern Child Health J.* 2013;17:661–666.
 37. Mashayekhi SO, Dilmaghanizadeh M, Sattari MR. A survey on the consumption, knowledge and attitude of pregnant women toward the effects of folic acid on pregnancy outcome in Tabriz. *Iran J Child Neurol.* 2011;5: 35–42.

38. Yajnik CS, Deshpande SS, Jackson AA, Refsum H, Rao S, Fisher DJ, et al. Vitamin B12 and folate concentrations during pregnancy and insulin resistance in the offspring: the Pune Maternal Nutrition Study. *Diabetologia*. 2008;5:29–38.

© 2021 Alanazi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/79482>