



Performance Characteristics and Economy of Using Noodle Waste as Replacement in the Diets of Cockerel Finisher Chickens

**A. J. Omole^{1*}, O. O. Obi¹, O. O. Adejinmi², F. O. Ajasin²
and J. A. Fayenuwo¹**

¹*Obafemi Awolowo University, Institute of Agricultural Research and Training Moor Plantation, Ibadan, Nigeria.*

²*Federal College of Animal Health and Production Technology, Ibadan, Nigeria.*

Authors' contributions

This work was carried out in collaboration between authors AJO and OOO initiated the study while all authors were involved literature searches, data collection and analyses, writing, reading and approving of the final manuscripts

Research Article

Received 27th April 2013
Accepted 25th June 2013
Published 25th September 2013

ABSTRACT

Noodle Waste (NW) is a by-product of food processing industry and it is cheaper in price compared to maize. The objective of this study was to determine the effect of replacing maize fraction of the diet of cockerel finisher chickens with NW on performance, cost benefits, blood haematology and serum biochemicals. One hundred and twenty cockerel finisher chickens of mean weight of 689.01 ± 2.4 g were randomly selected and allotted to 4 dietary treatments. Each treatment was replicated thrice with ten birds per replicate in a completely randomized design. Four diets were formulated to contain 0% (T₁), 50% (T₂), 75% (T₃) and 100% (T₄) of NW as replacement for maize fraction of the entire diet. Parameters measured or calculated were weight gain, feed intake, feed conversion ratio, total feed cost, and cost per weight gain, dressing percentages and blood indices such as Packed Cell Volume (PCV), Red Blood Cell (RBC) count and White Blood Cell (WBC) and total protein etc. Significant differences were observed in the mean total feed intake of the birds fed diet containing 0% and 100% NW as replacement for maize ($P=0.05$). The mean total weight gain of birds fed diet containing 100% NW and the control diet were significantly influenced by the dietary treatments ($P=0.05$). The feed conversion ratios were

*Corresponding author: Email: omoleboye@yahoo.com;

relatively similar ($P>.05$) in diet containing 0% and 100% NW as replacement for maize. The mean dressing percentage in all the treatments were relatively the same ($P>.05$). The dietary treatments had no significant effect on the PCV, RBC and WBC and blood total protein ($P>.05$). The lowest cost/weight gain was recorded in the diet containing 100% NW as replacement for maize. Based on the present results NW could replace maize fraction of the diet partially or wholly without any adverse effect on performance and health status but at reduced cost.

Keywords: Cockerel, cost/weight gain; energy; feed efficiency; dressing percentage; noodle waste; replacement.

1. INTRODUCTION

The animal protein intake of average Nigerian and African in general is low when compared to reports obtainable from Developed Nations of the world such as Great Britain, United States of America, Italy and Japan and this could be attributed to inadequate supply and high cost of animal sources of protein such as beef, pork and poultry products etc. [1,2,3]. There is need to improve livestock industries in Nigeria in order to meet the demand for meat and other livestock products. Unavailability of feed at affordable price constitutes a major limitation to the expansion of poultry and other livestock industries in Africa [4,5,6]. Conventional feed ingredients such as maize, soybean meal, groundnut cake meal and bone meal are expensive because of industrial uses and human consumption, hence the use of alternative feed resources must be encouraged in order to reduce feed cost which constitutes up to 60-70% of total cost of production. Noodle Waste (NW) is one of the alternative source of energy in the diet of livestock. According to Eniolorunda et al. [6] the protein content is about 9.56% with high metabolizable energy. The fat content is higher than that of maize. Moreover, the cost of NW is relatively cheaper than the cost of maize of the same quantity. It was concluded by Eniolorunda et al. [6] that maize could be partially replaced with NW without adverse effect on growth, egg and carcass quality of laying chicken. There is paucity of information on the use of NW in the diet of cockerel chicken hence, this study was designed to evaluate performance characteristics, carcass values, cost benefits and haematological indices of cockerel finisher chickens fed diet containing NW as partial or total replacement for maize in the entire diet.

2. MATERIALS AND METHODS

2.1 Source of the Test Ingredient and Experimental Design

A total of one hundred and twenty, 8 weeks cockerel chickens (Harco-Black) of mean weight $689.01 \pm 2.4g$ were used for the feeding trial. Four diets, T_1 , T_2 , T_3 and T_4 were formulated to contain 0, 50, 75 and 100% NW as replacement for maize in a growing-finishing cockerel diets respectively. NW was purchased from a feed mill in Ibadan, Oyo state, Nigeria. It was ground before incorporating with other feed ingredients. The diets were formulated to contain about 16% Crude Protein (CP) and energy of about 2800Kcal/kg ME (Table 1). Thirty birds each were randomly allotted to each of the treatment in a completely randomized design and each treatment was replicated thrice with 10 birds per replicate.

Table 1. Gross composition of the experimental diets

Ingredients (%)	Cost (N/kg)	T₁	T₂	T₃	T₄
Maize	62.00	40	20	10	0.0
Noodle waste	50.50	0	20	30	40
*Others fixed ingredients		60	60	60	60
Total		100	100	100	100
Cost/kg (N)		53.15	52.12	50.76	49.05
Calculated Composition					
Crude protein (%)		15.68	15.90	16.21	16.35
**Energy (Kcal ME/kg)		2836.2	2810.34	2798.	2768.34
				34	

*Others fixed ingredients: G.N.C., 10.0; Soybean meal, 9.2; Corn-bran, 25.3; Wheat offal, 8.0; Fish meal, 2; Bone meal, 1.5; Oyster shell, 3.5; Methionine, 0.1; Lysine, 0.1; Salt, 0.25; Premix, 0.25***.

**Metabolizable Energy calculated (KcalME /kg) = 37X %CP + 81 X%EE + 35.5 X % NFE [7].

***Vitamin-Mineral Premix (BIO-MIX) Supply the following per Kg: Vit.A, 5000IU; Vit.D3, 888.000IU; Vit.E, 12.000mg; Vit.K3, 1500mg; Vit.B1, 1000mg; Vit.B2, 2000mg; Vit.B6, 1600mg; Niacin, 12.000mg; Pantothenic acid, 2000mg; Biotin, 1000mg; Vit.B12, 3000mg; folic acid, 15000mg; Choline Chloride, 60.000mg; Manganese, 10.000mg; Iron, 15000mg; Zinc, 800mg; Copper, 400mg; Iodine, 80mg; cobalt, 40mg; Selenium, 8000mg

2.2 Data Collection

Feed and water were given *ad libitum*. Daily Feed Intake (FI) was calculated by deducting the left over feed from a weighed quantity of feed supplied daily. The birds were weighed at weekly intervals to determine the weekly and subsequently the daily weight gain. Feed Conversion Ratio (FCR) was calculated as the ratio of FI to weight gain. Record on mortality was also taken. The feed cost and cost per weight gain were calculated. The prevailing market prices of the feed ingredients at Ibadan, Oyo State, Nigeria where the feed ingredients were purchased at the time of study were used to calculate the feed cost/kg feed and the cost of the feed to produce a unit weight (N/kg weight gain). At the end of the 10th week, blood samples were collected from 6 birds in each experimental group (i.e. 2 birds per replicate) for the determination of the haematological and serum biochemical parameters. The birds randomly selected were fasted from 6.00 pm to 6.00 am and bled early in the morning to avoid temporary elevation of blood metabolites by feeding. Approximately 5ml of blood was collected from jugular vein of chickens into two sets of sterilized glass bottles containing Ethylene Diamine Tetra-Acetic Acid (EDTA) for haematology and without anticoagulant for serum separation. Samples for haematological study were collected into sample tubes containing EDTA as anticoagulant while serological samples were collected into sample tubes containing no EDTA. Serum was obtained after the blood was allowed to stand for 2h at room temperature and centrifuged at 2.000 revolutions per minute (rpm) for 10 minutes to separate the cells from the serum.

2.3 Blood and Feed Analyses

Haematological parameters included the Packed Cell Volume (PCV), Red Blood Cells (RBC), Haemoglobin (Hb) concentration, White Blood Cells (WBC), mean corpuscular volume, mean corpuscular haemoglobin, and mean corpuscular haemoglobin concentration. Total protein, albumin, globulin, and creatine formed the biochemical data were determined [8,9]. Blood samples were analyzed for haematological parameters according to routinely

available clinical methods [10]. The chemical composition of the test ingredient, experimental diets were carried out according to the method [11].

2.4 Data Analysis

All data were subjected to statistical analysis of variance and means were separated if found significantly different by Duncan multiple range test [12].

3. RESULTS AND DISCUSSION

The CP of NW and maize were numerically similar (Table 2) which was in agreement with the findings of [6]. The CP content of about 16% of the experimental diets was relatively similar to the report [13,14].

Table 2. Determined proximate composition of maize, noodle waste (NW) and experimental diets

Parameters	Maize	NW	T ₁	T ₂	T ₃	T ₄
Dry matter	93.78	92.74	95.96	94.67	94.87	95.45
Crude protein	9.89	10.02	15.56	15.59	15.69	15.76
Crude fibre	5.76	4.58	5.97	5.87	5.81	5.64
Ether extract	6.78	8.34	4.98	5.58	5.99	6.98
Ash	11.89	12.24	11.23	11.55	11.69	11.99
Nitrogen free extract	65.68	64.82	62.26	61.41	60.82	59.63

Significant differences were observed in the mean total FI of the birds fed diet containing 0% and 100% NW as replacement for maize ($P < .05$) (Table 3). The highest FI was recorded in T₄. There was no significant difference in the mean total FI in T₁, T₂ and T₃ ($P > .05$). High protein content of the test ingredient would have been responsible for improvement in numerical increased FI. The improvement in FI from T₁ to T₄ could also be due to decrease in the caloric level as the level of NW increased in the diet from 0% to 100% and the reports were in agreement with the findings of [3,6]. The mean total weight gain of birds fed diet containing 100% NW and the control diet were significantly influenced by the dietary treatments ($P < .05$). Birds in the control diet had higher weight than those in T₄ ($P < .05$). The FCR was relatively similar ($P > .05$) in diet containing 0%, 50, 75 and 100% NW as replacement for maize. The relative similarity in performance observed in the birds fed control diet and those fed 100% NW as replacement for maize implied that maize could be replaced partially or wholly by NW in the diet of cockerel finisher chicken without affecting the growth and feed efficiency. The low mortality recorded across the treatments may be an indication that NW do not present anti-metabolic compounds that could adversely affect health status of the birds. The cost per kg feed and total feed cost reduced as the level of NW in the diet increased from T₁ to T₄. The lowest cost/kg feed of N159.25 was recorded in diet containing 100% of NW as replacement for maize while the highest cost per weight gain of N166.15 was recorded (Table 3).

Table 3. Summary of performance characteristics and cost benefits of cockerel finishers chicken fed NW meal based diet as replacement for maize

Variables (Means)	T₁	T₂	T₃	T₄	± SEM
Initial weight. (g)	689.41	687.92	691.44	688.73	5.89
Final weight. (g)	2905.6 ^a	2907.4 ^a	2894.5 ^a	2884.3 ^b	7.45
Weight gain (g)	2216.19 ^a	2219.48 ^a	2213.06 ^a	2195.57 ^b	7.12
Total feed intake (g)	6936.68 ^a	6946.97 ^a	6949.0 ^a	7135.60 ^b	15.15
Feed conversion ratio	3.13	3.13	3.14	3.25	0.23
Mortality (%)	3.33	0.00	3.33	3.33	0.03
Cost/kg feed (₦ /kg)	53.15 ^a	52.12 ^a	50.76 ^{ab}	49.05 ^b	2.05
Total feed cost (₦ /kg)	368.86 ^a	362.23 ^b	352.78 ^c	349.73 ^c	4.78
Cost/wt. gain (₦ /kg)	166.15 ^a	163.16 ^{ab}	159.62 ^b	159.25 ^b	3.05

Means with different superscripts along the same row are significantly different (P<.05)

The mean dressing percentages in all the treatments were relatively the same as shown in Table 4. The values ranged between 72.88% and 73.42%, similar to the report of (13, 14). The heart, lung and liver weights were not affected by the inclusion of noodle as replacement for maize (Table 4). There was no depressive effect of NW on PCV as the level increased from 0-100% as replacement for maize in the diet (P>.05). The PCV values fell within the recommended values according to Afolabi et al. [15].

Table 4. Carcass composition of cockerel finishers chicken fed NW meal based diet as replacement for maize

Variables (Means)	T₁	T₂	T₃	T₄	± SEM
Live weight (g)	2901.6 ^a	2903.4	2891.3 ^a	2876.3 ^b	6.21
Eviscerated weight(g)	2130.07 ^a	2131.68 ^a	2108.9 ^b	2096.2 ^b	11.89
Dressing percent (%)	73.41	73.42	72.94	72.88	3.43
Heart weight (%)	0.65	0.64	0.65	0.63	0.11
Liver weight (%)	2.68	2.65	2.68	2.62	0.23
Lung weight (%)	1.79	1.76	1.77	1.75	0.13

Means with different superscripts along the same row are significantly different (P<.05)

The Hb concentration of the birds was not significantly affected by the dietary inclusion level of NW in the diet of growing-finishing cockerel (Table 5). The values of Hb count ranged between 11.33 and 11.44% (P<.05) and this indicates that the oxygen carrying capacity of the bird's blood by Hb was relatively the same. The WBC counts were relatively the same across the treatments and the values obtained were in agreement with the recommended values of cockerel finisher chickens [9,16]. The WBC plays a major role in defending the body against disease-producing bacteria, viruses and fungi, a deficiency in WBC may result in an increased susceptibility to infections. A decrease in WBC counts is a reflection of the decline in the production of WBC for defensive action against infection and it has also been established that an animal with decrease in the level of Hb in the blood is indicative of poor nutrition including dietary deficiency of iron, amino acid and vitamins [9,15,17]. The results obtained in the study also buttress the fact that NW had no detrimental effect on health status of the birds and could be used as alternative feed resource in the diet of cockerel finisher chickens.

Table 5. Haematology parameters of cockerel finishers chicken fed NW meal based diet as replacement for maize

Variables (Means)	T₁	T₂	T₃	T₄	± SEM
PCV (%)	27.69	28.45	28.91	27.89	2.23
R.BC(x 10 ⁶ /μ1)	4.21	4.20	4.24	4.33	0.67
Haemoglobin (g/dl)	11.35	11.33	11.43	11.44	1.43
WBC (x 10 ³ /μ1)	11.98	11.98	11.86	12.01	1.76
MCHC%	32.46	32.32	32.45	33.01	2.43
Neutrophil %	34.67 ^b	35.43	34.97	34.88	3.12
Monocyte%	0.41	0.41	0.43	0.45	0.11
Eosinophil	0.14	0.13	0.13	0.15	0.05

Means along rows with different superscript are significantly different from each other (P<.05)

The results of serum biochemical values presented in Table 6 shows that the total protein, albumin and globulin of the birds in all the treatments were not significantly influenced by the inclusion of NW in the diets (P<.05). Numerically, birds fed diet 4 had slight higher total protein and albumin levels than those fed 0, 50 and 75% NW as replacement for maize and this could be due to high inclusion levels of NW and the report is in agreement with the findings of Iyayi and Tewe [18] who reported that total protein contents of the blood depend on the quantity and quality of the protein supplied in the diet. Based on the present results of FCR, dressing percentages, cost per weight gain and blood indices, NW could replace maize fraction of the diet partially or wholly without any adverse effect on health status but at reduced cost.

Table 6. Serum biochemical parameters of cockerel finishers chicken fed NW meal based diet as replacement for maize

Parameters (Means)	T₁	T₂	T₃	T₄	± SEM
Total protein (g/dl)	5.58 ^b	5.59	5.61	5.77	0.68
Globulin (g/dl)	3.34	3.36	2.37	3.42	0.36
Glucose Mg/dl)	156.8	158.23	160.3	161.34	6.34
Albumin g/dl	2.61	2.63	2.65	2.70	0.12
Creatine Mg/dl	1.21	1.23	1.30	1.33	0.05

Means along rows with different superscript are significantly different from each other (P<.05)

4. CONCLUSION AND RECOMMENDATION

Based on the present results NW could replace maize fraction of the diet partially or wholly without any adverse effect on performance and health status but at reduced cost. It could be recommended that farmers should include NW in the diet of cockerel finisher chickens for cost reduction.

ACKNOWLEDGEMENT

We wish to thank the management and all members of Staff of The Institute of Agricultural Research and Training Communities, Moor Plantation, Ibadan for their encouragement towards the success of the article. Our sincere appreciation also goes to Mr. E.S.O. Arabambi, Mrs. O. Ajayi (Mama Ibeji), Mrs. A. Fatoyibo for the typing of the manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Longe GO. Replacement value of biscuit waste for maize in broiler diets. *Nig. J. Anim. Prod.* 1987;13(1-2):70-78.
2. FAO. Food and Agricultural Organization of the United Nations. *The State of the World's Animal Genetic Resources for Food and Agriculture*. Rischkowasky and D. Pilling (Eds.) FAO, Rome, Italy; 2007.
3. Olayeni TD, Farinu GO, Ojebiyi OO. Replacement Value of biscuit waste on the performance and egg quality parameters of laying hens. *Proc. 32nd Ann.* 2007;313-230.
4. Obioha FC. *Guide to Poultry Production in the tropics*. Acena publishers, Enugu, Nigeria. 1992;74.
5. Atteh JO. Principles and practice of livestock feed manufacturing. Adlek Printers, Ilorin, Nigeria. 2002;52-58.
6. Eniolorunda OO, Taiwo BBA, Oyewumi OO, Adeyemi OA. Performance of laying hens fed graded levels of Indomie waste as replacement for maize in a humid tropical environment. *Res. J. Anim. Sci.* 2008;2(5):135-138.
7. Pazuenga U. Feeding parent stock. *Zootecnica International*. 1985;22-24.
8. Frandson RD. *Anatomy and Physiology of farm animals*. Lea Febinger Publications, Philadelphia, 1st Edition. 1981;86-94.
9. Mitruka HM, Rawnsley SK. *Clinical, Biochemical and Haematology reference in normal experimental animal*. Mason, NY. 1997;287-380.
10. Scalm OW, Jane NC, Carol EA. *Veterinary Haematology*, 3rd edition Lea and Febigar, Philadelphia, U.S.A. 1975;160-210.
11. Association of Official Analytical Chemist (A.O.A.C.). *Official Methods of Analysis*, 13th Edition, Washington, D.C. ; 1990.
12. SAS Statistical Analysis System SAS Stat. Version 9, SAS Institute Inc. Gary, NC 27513 USA; 2000.
13. Fapohunda JB, Adejinmi OO, Okpeze CN, Owosibo OA, Bamigbose AM. Effect of replacing maize with biscuit waste in the diet of cockerels Proceeding of the 13th Annual conference of the Animal Science Sept. 15-19, 2008, ABU, Zaria. 2008;318-321.
14. Adejinmi OO, OkpezeCN, Obi OO, Omole AJ, Kehinde AS, Awe OA. The use of chicken egg shell as an alternative source of calcium in the diet of cockerel chicken. *Journal of Agriculture and Social Sciences*. 2011;34-38.
15. Afolabi O, Oladimeji H. Haematological studies of some avian species. *International Journal of Poultry Science*. 2003;30(2):24.
16. Lamb GN. *Manual of Veterinary Laboratory Technique*. CIBA-GEIGY, Kenya. 1981; 70-110.
17. Aletor VA, Egberongbe O. Feeding differently processed soybean, Part 2 and assessment of haematological responses in chicken. *Die Nahrung*. 1992;36:367-370.

18. Iyayi EA, Tewe OO. Serum total protein, urea and creatine levels as indices of quality cassava diets for pigs. *Tropical Veterinary*. 1998;16:57-67.

© 2013 Omole et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<http://www.sciencedomain.org/review-history.php?iid=236&id=2&aid=2051>