



**Bionature**

Volume 44, Issue 1, Page 11-15, 2024; Article no.BN.1514  
ISSN: 0970-9835 (P), 0974-4282 (O)

# Exploring the Influence of Mucin on Lipid Bilayers: Insights into Phase-Dependent Interactions

**Jin-WonPark<sup>a++\*</sup>**

<sup>a</sup> Department of Chemical and Biomolecular Engineering, School of Energy and Biotechnol, Seoul National University of Science and Technol, South Korea.

## Author's contribution

The sole author designed, analyzed, interpreted and prepared the manuscript.

## Article Information

DOI: 10.56557/BN/2024/v44i11983

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://prh.globalpresshub.com/review-history/1514>

**Short Communication**

**Received: 16/01/2024**

**Accepted: 19/03/2024**

**Published: 26/03/2024**

## ABSTRACT

Changes in the behavior of the prepared lipid bilayer according to each phase of the layer, induced by mucin incorporation, were investigated. 8-Aminonaphtalene-1,3,6-trisulfonic acid disodium salt (ANTS) was encapsulated only in the water-soluble interior of the vesicle. *p*-xylene-bis(*N*-pyridinium bromide) (DPX) settled only outside of the vesicles to quench the ANTS. The calibration was established for the fluorescence with the encapsulated ANTS separated from the DPX (100%) and the ANTS mixed with DPX (0%). The changes were found due to the layer phase over the ratio of mucin to lipid. The changes are determined by the stability of the vesicles.

**Keywords:** Mucin; lipid bilayer; vesicle; fluorescence; layer phase.

## 1. INTRODUCTION

Mucin is commonly observed on luminal surface of animals [1]. This group consists of a linear

peptide with oligosaccharide [2,3]. Additionally, a hydrophobic transmembrane stretch of the mucin is used to its anchorage to the plasma membrane and signaling [1]. The mucin has

<sup>++</sup> Professor and Chair;

<sup>\*</sup>Corresponding author: Email: [jwpark@seoultech.ac.kr](mailto:jwpark@seoultech.ac.kr);

been recognized as being involved in the development of cancer, especially adenocarcinomas [4]. Furthermore, its association with tumor has been identified [5].

The mucin has been believed to limit the lateral movement of the lipids, because the mucin was distributed at the interface of the lipid layer [6,7]. The mucin became more stable due to the presence of lipid [8]. Particularly, the phosphatidylcholine was found to have an affinity for mucin [9]. Additionally, the phosphatidylcholine was observed to penetrate more easily into the mucin compared to other nanocarriers [10].

Little has been found for the effect of mucin on the layers. Although the behaviors of lipid monolayers were investigated with the addition of a compound, a bilayer is more complicated than an assembly of two monolayers [11-15]. In this study, the purpose is to investigate the effect of mucin on lipid bilayers made of each-phase layer.

## 2. MATERIALS AND METHODS

### 2.1 Vesicle Preparation

Ten mg of dipalmitoylphosphatidic acid (DPPA) or dioleoylphosphatidic acid(DOPA) from Merck (Rahway, NJ) was dissolved in 2 ml of tert-butyl methyl ether. To this solution was added 100  $\mu$ l DI water of 25 mM 8-aminonaphthalene-1,3,6-trisulfonic acid disodium salt(ANTS), 10 mM Tris-HCl at pH 7.4. For the formation of the desired inverse-micelles, the extrusion of the above solution was performed through the 50 nm pores


of 78 mm diameter PTFE membranes at a temperature higher than the transition of the chosen lipid. To the 10 ml Tris-HCl buffer of 90 mM p-Xylene-bis(N-pyridinium bromide)(DPX), either dioleoylphosphatidylcholine (DOPC) or dipalmitoylphosphatidylcholine (DPPC) was added continuously in a drop way less than 10  $\mu$ l micelle-solution. This solution underwent through the centrifugation (3700  $\times$  g), and its supernatant only was the vesicle solution [16].

### 2.2 Vesicle Confirmation

From the diameters, both micelles and vesicles were confirmed using dynamic light scattering. As shown in Fig. 1, each diameter was  $75 \pm 10$  and  $80 \pm 10$  nm. The refractive index and the viscosity of the solvent are 1.3686 and 0.23 cP [17]. In addition, little change in the 530 nm fluorescence intensity with pH change outside vesicle indicated that each layer was secured.

### 2.3 Fluorescence Measurement

The fluorescence intensity was monitored at the desired ratio of mucin to PC using a fluorometer (Perkin-Elmer, Waltham, MA). The intensity change generated by the mucin means that the layer arrangement was disturbed because the change in the ANTS fluorescence intensity is caused only due to the mixing of ANTS and DPX. The calibration was established for the fluorescence with the encapsulated ANTS separated from the DPX (100%) and the ANTS mixed with DPX (0%) [18]. The quantitative intensity was found to understand the mucin effect.



**Fig. 1. Diameter data of (A) micelles and (B) vesicles**

### 3. RESULTS AND DISCUSSION

#### 3.1 Vesicle Fusion Induced by Mucin

The results for each phase of the layer are summarized with respect to the mucin ratio (Table 1). Each phase corresponded to the intensity. For the solid outer, the intensity dropped to about 0.5 ratio. In addition, the change in the lipid phase was generated by the cosmotropic effect of mucin on the water around the headgroup, because most lipids were associated with the mucin at the ratio. The change was saturated at 0.5 ratio.

However, for the liquid outer, the difference in the intensity rose. Little change was observed for any phase of the inner in the case where the outer was liquid. In other words, the mucin-caused-induction of the vesicle fusion was generated only at the solid outer. Therefore, the intensity change was determined dominantly by the outer phase due to the layer exposure to the mucin. In addition, the influence of difference in the outer phase was considered under the identical phase of the inner. The behavior of the

intensity was not identical each other (Fig. 2). In the case of the liquid inner, the lowest intensity was identical for any phase of the outer (Fig. 2 (A)). However, in the solid inner, the intensity of the solid outer became from higher to lower rather than the liquid outer (Fig. 2 (B)). These results indicated that mucin, only outside the vesicle, disturbed both outer solid phase and inner solid phase. The liquid phase of lipid layer has more flexibility to cushion the disturbance because its arrangement is less organized than the solid. In other words, the mucin little disturbed both liquid layers. The previous investigation was consistent with this interpretation [19].

#### 3.2 Effect of Mucin on Vesicle Inner


The intensity of the solid outer was expected similar with that of the liquid outer because the mucin induced the fluidity on the solid layer. However, the observation was different. The discrepancy from the expectation seemed to be caused by the tail-group mismatch between each layer. The cosmotropic effect increased the distance between the head-groups of the

**Table 1. Fluorescence intensity(FI) for each layer phase after the mucin addition**


	Solid Outer											
	Solid Inner						Liquid Inner					
	Ratio of Mucin to Lipid						Ratio of Mucin to Lipid					
FI (%)	0	0.1	0.3	0.5	0.7	1.0	0	0.1	0.3	0.5	0.7	1.0
	100	90	70	50	50	50	70	60	45	30	30	30

	Liquid Outer											
	Solid Inner						Liquid Inner					
	Ratio of Mucin to Lipid						Ratio of Mucin to Lipid					
FI (%)	0	0.1	0.3	0.5	0.7	1.0	0	0.1	0.3	0.5	0.7	1.0
	70	70	70	70	70	70	30	30	30	30	30	30



**Fig. 2. Fluorescence intensity of vesicles with solid outer (---) and with liquid outer (—) at 0.5 ratio. (A) liquid inner and (B) solid inner**



**Fig. 3. Mucin effect on lipid layer. Outer layer is exposed to mucin. (A) Both solids, (B) Solid outer-layer, (C) Liquid outer-layer, and (D) Both liquids**

solid outer. Although the lipids of the solid outer were far from the others, little fluidity was in the tail-groups. Therefore, the mucin disrupted the arrangement of the layer tail-groups. If the tail-group was liquid, the layer was less rigid over the disruption. This interpretation corresponded to the lower intensity of the solid outer compared to the liquid outer in the case of the solid inner, as shown in Fig. 3 (next page). The effect induced by mucin was almost identical with that by hydroxybutyric acid, although the mucin was much larger than the hydroxybutyric acid. This fact may imply that only a part of the mucin affected the lipid layer [18].

From about 20% reduction in the fluorescence intensity, it has been inferred that the change in the structure occurred to most of the vesicles. All results found out in this study may attribute to the cosmotropic and volumetric effects, which disrupt the head-group packing and the tail-group arranging. Since the liquid of layers is critical for the functions, the liquefaction by the mucin may be related to the mechanism.

#### 4. CONCLUSION

The mucin-induced vesicle instability was monitored according to layer phase using the fluorescence intensity. The intensity in the solid outer was inversely proportional to the mucin ratio up to 0.5. In the liquid, the behavior of the intensity was distinguished from the solid. These results may attribute to the cosmotropic and volumetric effects, which disrupt the head-group packing and the tail-group arranging. The mucin study may be related to the biological mechanism connected to cellular processes.

#### COMPETING INTERESTS

Author has declared that no competing interests exist.

#### REFERENCES

1. Marin F, Luquet G, Marie B, Medakovic D, Molluscan shell proteins: Primary structure, origin, and evolution. *Curr. Top. Dev. Biol.* 2007;80:209–276. Available: [https://doi.org/10.1016/S0070-2153\(07\)80006-8](https://doi.org/10.1016/S0070-2153(07)80006-8)
2. Strous GJ, Dekker J, Mucin-type glycoproteins. *Crit. Rev. Biochem. Mol.* 1992;27:57–92. Available: <https://doi.org/10.3109/10409239209082559>
3. Shi L, Caldwell KD. Mucin adsorption to hydrophobic surface. *J. Colloid Interface Sci.* 2000;224:372–381. Available: <https://doi.org/10.1006/jcis.2000.6724>
4. Maill AS, Analysis of mucins: Role in laboratory diagnosis. *J. Clin. Pathol.* 2008;61:10118–11024. Available: <https://doi.org/10.1136/jcp.2008.058057>
5. Bansil R, Turner BS, The biology of mucus: Composition, synthesis and organization. *Adv. Drug Deliv. Rev.* 2018;124:3-15. Available: <https://doi.org/10.1016/j.addr.2017.09.023>
6. Wiedmann TA, Herrington H, Deyea C, Kallick D. Analysis of the diffusion of bile salt/phospholipid micelles in rat intestinal mucin. *Chem. Phys. Lipids.* 2001;112:81–92. Available: [https://doi.org/10.1016/s0009-3084\(01\)00168-2](https://doi.org/10.1016/s0009-3084(01)00168-2)
7. Millar TJ, Tragoulias ST, Anderton PJ, Ball MS, Miano F, Dennis GR, Mudgil P. The surface activity of purified ocular mucin at the air-liquid interface and interactions with meibomian lipids. *Cornea.* 2006;25:91–100. Available: <https://doi.org/10.1097/01.ico.0000164779.87795.3c>

8. Rogunova MA, Blackwell J, Jamieson AM, Pasumar-Thy M, Gerken TA. Effect of lipids on the structure and rheology of gels formed by canine submaxillary mucin. *Biorheology*. 1997;34:295–308. Available:[https://doi.org/10.1016/s0006-355x\(98\)00006-7](https://doi.org/10.1016/s0006-355x(98)00006-7)
9. Korytowski A, Abuillan W, Amadei F, Makkya A, Gumiero A, Sinning I, Gauss A, Stremmel W, Tanak, M. Accumulation of phosphatidylcholine on gut mucosal surface is not dominated by electrostatic interactions. *Biochim. Biophys. Acta (BBA)-Biomembr*. 2017;1859:959–965. Available:<https://doi.org/10.1016/j.bbamem.2017.02.008>
10. Alp G, Aydogan N, Lipid-based mucus penetrating nanoparticles and their biophysical interactions with pulmonary mucus layer. *Eur. J. Pharm. Biopharm*. 2020;149:45–57. Available:<https://doi.org/10.1016/j.ejpb.2020.01.017>
11. Crowe JH, Whittam MA, Chapman D, Crowe LM. Interactions of phospholipid monolayers with carbohydrates. *Biochim. Biophys. Acta*. 1984;769:151-159. Available:[https://doi.org/10.1016/0005-2736\(84\)90018-x](https://doi.org/10.1016/0005-2736(84)90018-x)
12. Diaz S, Lairi3n F, Arroyo J, Biondi de Lopez AC, Disalvo EA. Contribution of phosphate groups to the dipole potential of dimyristoylphosphatidylcholine membranes. *Langmuir*. 2001;17:852-855. Available:<https://doi.org/10.1021/la000683w>
13. del Luzardo M, Amalfa CF, N3nez AM, Diaz S, Biondi de Lopez AC, Disalvo EA. Effect of trehalose and sucrose on the hydration and dipole potential of lipid bilayers. *Biophys. J*. 2000;78:2452-2458. Available:[https://doi.org/10.1016%2Fs0006-3495\(00\)76789-0](https://doi.org/10.1016%2Fs0006-3495(00)76789-0)
14. Nagle JF. Theory of lipid monolayer and bilayer chain-melting phase transitions. *Faraday Discuss. Chem. Soc.* 1986;81: 151-162. Available:<https://doi.org/10.1039/DC9868100151>
15. Nagle JF, Tristram-Nagle S, Structure of lipid bilayers. *Biochim. Biophys. Acta*. 2000;1469:159-195. Available:[https://doi.org/10.1016%2Fs0304-4157\(00\)00016-2](https://doi.org/10.1016%2Fs0304-4157(00)00016-2)
16. New RRC. *Liposomes: A practical approach*, Academic Press, New York, USA; 1990:20-41.
17. Lide DR. *CRC handbook of chemistry and physics: A ready-reference book of chemical and physical data*, 85th ed., CRC Press, New York, USA. 2005;220.
18. Lee G, Park J-W. Effect of hydroxybutyric-acid on lipid bilayers with respect to layer phase. *Journal of the Korean Applied Science and Technology*. 2022;39:720-726. Available:<http://dx.doi.org/10.12925/jkocs.2022.39.5.720>
19. Park J-W. First-leaflet phase effect on properties of phospholipid bilayer formed through vesicle adsorption on LB monolayer. *J. Membr. Biol*. 2010;237:107-114. Available:<https://doi.org/10.1007/s00232-010-9311-0>

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://prh.globalpresshub.com/review-history/1514>