



A Scoping Review of the Genotypic Diversity of *Streptococcus mutans*

**K. Kunguma Supraja^{a≡∅}, Saravanan Poorni^{a,b≡‡#*}, Nivedhitha^{c≡†},
Manali Ramakrishnan Srinivasan^{a≡†}, M. Jeevitha^d and Selvaraj Jeyaraman^e**

^a Department of Conservative Dentistry and Endodontics, Sri Venkateswara Dental College and Hospital, Chennai, India.

^b Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences, Saveetha University, India.

^c Department of Conservative Dentistry and Endodontics, Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai - 600077, Tamil Nadu, India.

^d Department of Periodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, India.

^e Department of Biochemistry, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i62B35175

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/82438>

Review Article

**Received 20 October 2021
Accepted 26 December 2021
Published 29 December 2021**

ABSTRACT

Aim: This paper reviews the literature on the pathogenic genotypes of *S. mutans* that may be more virulent colonizers and the phenotypic variability of its main virulence factors.

Methods: A thorough literature search on *S. mutans* was performed and the relevant data supporting its association to dental caries were extracted.

[≡] Dr.;

[∅] Post Graduate Student;

[‡] PhD Scholar;

[#] Professor;

[†] Professor and Head;

*Corresponding author: E-mail: poornii@yahoo.com;

Results: Dental caries is a microbial disease caused by frequent intake of dietary sugars. Fermentation of sugars by biofilm microbiota produces acids that disrupt microbial homeostasis and cause dissolution of tooth minerals. Identifying *S. mutans* as the most important cariogenic microorganism has led to the design of target specific preventive measures that intend to reduce its presence in oral cavity.

Conclusion: Due to environmental changes and selective pressure in the oral cavity, *Streptococcus mutans* endure extensive genotypic diversity thereby exhibiting new physiological and metabolic properties. However, the role of the variants are poorly understood.

Keywords: Dental caries; genotypic diversity; *Streptococcus mutans*.

1. INTRODUCTION

Dental caries is a ubiquitous chronic diseases affecting people all through their lives. It is a complex multifactorial microbial disease affecting billions of people globally, thereby posing a major public health concern. Although diverse factors aid in the development of dental caries, the disease is mainly driven by the resident cariogenic microbiota which produces acids by metabolizing the dietary carbohydrates. The production of such acids and maintenance of its low pH over prolonged duration in the biofilm of the microbiota, present in proximity to the teeth, predisposes to formation of dental caries by dissolution of the tooth minerals, in an opportunistic manner. Currently the models of dental caries etiology focuses on relating this disease to a microbial ecological shift, which is based on a physiologic imbalance between tooth mineral and biofilm [1,2]. While dental caries is not a typical infectious disease it is important to know about the infectious and transmissible characteristics of the endogenous oral microflora.

Amongst the plethora of microorganisms present in the oral cavity, the Mutans streptococci have been implicated as the primary agent related to dental caries initiation in humans [3-5]. By virtue of its contribution to the formation of the dental biofilm matrix, its capacity to produce large quantities of organic acids, and its propensity to outcompete non-cariogenic commensal species at low pH conditions [6]. Decades of research on Mutans Streptococci association to dental caries has conclusively narrowed down on to *S. mutans*. However *S. mutans* can be isolated from individuals either with or without a history of caries [7,8]. Thus existence of variations among the colonizing *S. mutans* is the reason for the subsequent development and progression of caries among different individuals. To understand the infectious nature of dental caries it may be useful to study the genotypes of *S. mutans* to appreciate its diversity. According to the official

Guidelines for Human Gene Nomenclature, a gene is defined as a DNA segment that contributes to phenotype/function [9]. Genotype is the genetic constitution of a cell, an individual or an organism. Also gene regulation is found to play a vital role in the control of the expression of virulence factors such as synthesis of polymeric substances, adhesion, acidogenicity, aciduricity and variability. Diversity is the number of different *S. mutans* genotypes found within an individual. Genetic diversity of *S. mutans* remains as a topic of interest, as this provides an explanation for the contrasting caries status in people harbouring it [10]. Thorough understanding of these microbial species and their colonisation stratagems may help in the diagnosis, risk assessment, development of new treatment strategies for caries, so as to prevent disease and promote health in addition to standard prevention treatments.

Human oral microflora is composed of an abundant mix of microorganisms that are part of a multispecies biofilm complex. The oral cavity being a dynamic environment allows for the selection of certain organisms to not only adapt metabolically but also to develop its potential and genomic content in the biofilm [11]. Although microorganisms belonging to the same genus and species present a common gene set denoted core-genome, which is essential for cellular functioning and the survival of the species, they can differ in their physiological and virulent properties because of their strain specific genes [12-15]. The rest of the genome, referred to the accessory functions, is generally known as the not essential or dispensable genome and is not shared by all strains. Comparative genomic analysis between multiple genomes of individual species has revealed an extensive intra-species genomic diversity. The dispensable genome contributes to the diversity of the species and probably provides functions that are not essential for survival. It also confers selective advantages such as survival and adaptation to different

ecological niches, antibiotic resistance, and the capacity to colonize new hosts [14]. This diversity is generated by a variety of processes which include genome rearrangements, horizontal gene transfer by natural genetic transformation, and exposition to genetic mobile elements such as bacteriophages, plasmids, insertion elements, transposons, and genomic islands [12,16]. With the advent of new molecular tools over the past few decades there has been a substantial change in the understanding of dental caries microbial pathogenicity relative to bacterial detection and genotyping.

Rationale: The dental biofilm is composed of complex bacterial community, specific strains of *S. mutans* have to compete with other strains to establish colonization [17]. Gene regulation resulting in differences in genetic content of each *S. mutans* strain will also influence exhibition of phenotypic variability between the strains [18-20]. Knowledge of the various genotype of *S. mutans* with varied virulence factors and their correlation with other species is fundamental to understand their colonization and establishment in the same individual [21]. It has been suggested that not all strains have the same virulence capacity to promote the formation of dental caries [20,22]. Therefore, it seems reasonable to expect that at least some of the genes related to virulence are not distributed in the same way in the strains according to their caries status [23].

2. *S. mutans* SEROTYPES

S. mutans belong to the Mutans streptococci group along with nineteen other species [24]. It can be further classified into specific serotypes based on the cell wall antigenic polysaccharides known as rhamnose-glucose polymers [25,26]. *S. mutans* is found to have the highest genetic variation than other species present in humans and is classified into four serotypes (c, e, f, and k) [18]. Use of polymerase chain reaction (PCR) with primers designed based on the sequence differences of the *rgp* gene [18,27], has facilitated in determining the frequencies and distribution of these serotype in clinical isolates. Studies have revealed that serotype c is the frequent one corresponding to approximately 70-80% of the isolates in the oral cavity, followed by serotype e (20%), and serotypes f and k corresponding to less than 5% and 2% respectively [18,28-30]. This distribution also highlights different pathogenic patterns in *S. mutans*.

Environmental changes, selective pressures, and the presence of a variable genome in *S. mutans* cues its adaptation as well as strongly influence the acquisition of new physiological and metabolic properties that alter the dental biofilm homeostasis, leading to development of dental caries. Natural genetic transformation is a genetically programmed process which offers the recipient microorganism with the ability to acquire new phenotypic traits by facilitating the integration of dispensable genes [21,31]. Resulting in generation of a wide range of genome heterogeneity thereby promoting the emergence of resistance, genetic variation, and the rapid evolution of virulence factors [14,21,31,32]. This process is carried out as long as the cells have the ability to enter a physiological state known as competence which allows them to take up exogenous DNA from their environment and incorporate it into their genome [33]. It is a mechanism of ecological importance as *S. mutans* manages to adapt to the changing environments, originating from different selective pressures [21,34].

Although the existence of heterogeneity of *S. mutans* is widely known, very little is known about the mechanisms for the expression of pathogenic properties in specific genotypes. The employment of the state of the art molecular biology techniques in the various studies on *S. mutans* has provided enormous information on the evolution, pathogenesis, diversity, metabolic activities, and virulence properties of this species [35]. The three main natural strategies that generate genetic variations are:

- (1) Small local changes in the nucleotide sequence of the genome, which explains the presence of four serotypes
- (2) Intragenomic reorganization of segments of genomic sequences, and
- (3) Acquisition of DNA sequences from another organism

3. INTRAGENOMIC REARRANGEMENTS

Genomic rearrangements are mutations that change the gene content of a genome or the arrangement of the genes in a genome. They can be categorized as deletions, duplications, insertions, inversions, and translocations. They occur due to break in long stretches of DNA involving at least two different locations, followed by a re-ligation of the broken ends to produce a new chromosomal arrangement [36]. Sometimes this rearrangement could encompass a cluster of

genes depending on the size of the rearranged fragment [37]. Rearrangements therein influence the structure of the chromosome through disruption of an existing gene or by creation of a new gene, thereby affecting its expression instigating phenotypic variability [36].

In 2002 the complete genome sequencing of the *S. mutans* strain UA159 has helped unriddle the complexity and genetic specificity of *S. mutans* [21]. Since then UA159 is frequently used as a reference strain for variability studies [13]. Till date, a large number of *S. mutans* genomes have been sequenced and 188 genomic sequences of *S. mutans* are available in the NCBI database for the researchers. In comparative genomics studies the genomic sequences are often used to identify the virulence factors. It involves comparing genomes from different strains, then detecting the differences and similarities, thereby revealing common molecular and pathogenic mechanisms which can be related to specific phenotypic characteristics [38].

A pioneering study by Maruyama et al on *S. mutans* using comparative genomic analysis provided extensive information on the species-specific genetic content of *S. mutans*, while comparing strain UA159 with strain NN2025. The findings of the study were, although these strains belonged to the same serological group (serotype c) with expression of the same biochemical, adhesive, and cariogenic properties as well as similarity in 90% of their genes the strains presented eight variable regions with more than 30 genome rearrangements. The NN2025 strain genome contained eight strain-specific regions while the UA159 strain genome contained nine strain-specific regions [34].

Shields et al. utilized transposon sequencing technology and discovered that only 11% of the *S. mutans* UA159 genome is essential, with the presence of genes encoding products required for replication, translation, cell wall biogenesis, and the lipid metabolism, as well as genes necessary for survival, growth, and persistence colonization in both experimental and clinical conditions. It was suggested that *S. mutans* strains evolve through a process of chromosomal shuffling, which plays an important role in the genomic diversity of this species [32].

Song et al. sequenced the genomes of six *S. mutans* clinical isolates and compared them to the reference strains UA159 and NN2025,

focusing on characteristics related to pathogenicity. Genome alignment study revealed existence of chromosomal rearrangements amongst the strains. High variations in virulence-related genes between the strains was apparent. In addition to these findings the genomic regions required for *S. mutans* survival in different environments were also discovered [31].

4. HORIZONTAL GENE TRANSFER (HGT)

It is the movement of genetic information within a wide range of the bacteria which inhabit the human oral cavity [11]. As a result of it strain variants can be spawned due to loss, duplication, or modification of existing genes [12]. Often through this mechanism new phenotypic traits are acquired to provide a selective advantage to the microorganism. Therefore, due to the genomic diversity present among different isolates, the genome content of a single strain does not necessarily represent the genomic potential of certain species [34].

Hoshino et al. and Argimón et al. determined the origin of glycosyltransferases (GTFs) in the *Streptococcus* genus. This enzyme catalyses the synthesis of glucans from sucrose and are encoded by *gtf* genes. The authors proposed that the *Streptococcus* acquired the genes *gtf* by HGT and then were capable of forming cariogenic biofilms [39,40].

Cornejo et al. sequenced 57 *S. mutans* isolates to determine the general structure and the potential adaptive characteristics of both the core and dispensable genomes. The genomes of the studied strains were highly variable and their global genetic compositions were found to differ markedly from one isolate to another owing to the high HGT rate [15].

Meng et al. performed a pan-genome analysis of 183 *S. mutans* strains and determined that this species has an open pan-genome, thereby indicating that new genes can be found as more genomes are sequenced. An open pan-genome has been found to be associated with the adaptation of the bacterial species [38]. In addition, this is a distinctive characteristic of species which colonize diverse habitats and coexist with other microorganisms in large communities. It is typical of species that have the ability to exchange genetic material and have a high rate of HGT [41]. Hence, these findings suggest that the genome of *S. mutans* may be expanding gradually over time.

5. GENE REGULATION

S. mutans have been studied in some detail at genetic level, however many of the discovered genes have not been studied in the context of genetic competence until the past few years [42]. This can actually help us in understanding the cell-cell communication in *S. mutans* [43]. The various mechanisms of gene regulation in *S. mutans* include competence, signal transduction, quorum sensing, and small RNA (sRNA) regulation. For survival in a dynamic environment such as oral cavity, microorganisms were found to use regulatory systems to detect and respond rapidly to the stimuli generated [44]. Presentation of these regulatory genes involves complex coordinated processes that depends on the internal and external signals [45].

Shields et al. detected and characterized 20 novel genes that have a substantial impact on competition regulation and competence-related phenotypes. It has been studied that acquisition of natural competence is mostly transient and the genes encoding are not expressed as well [44]. The activation of the transcription of these genes take place in reaction to specific signals, and only when the environmental conditions are conducive. It is found out that in *S. mutans*, a network of genes and at least two peptide signaling molecules are accountable for the development of its genetic competence and its expression varies among the clinical isolates [31,40,46]. This was verified by Palmer et al., when they revealed that 15 isolates from geographically diverse patients with caries presented difference in the genetic content and phenotypic characteristics associated with virulence [47].

Strain UA159 were found to be naturally competent, transformable, and have a system committed to competence and quorum sensing, which is regulated by special signal peptides. Additionally, this systems also coordinates the development of antagonistic interactions and the synthesis of antimicrobial bacteriocins as a means to acquire the transforming DNA by competing with others. This is achieved either by eliminating closely related streptococcal species or through an altruistic suicide mechanism among a subpopulation of competent cells within the community of *S. mutans* [47-49].

S. mutans have also been found to be able to trigger a competence cascade. Besides the existing association between natural genetic

competence and stress response pathways, multiple environmental factors of the oral cavity can provoke a direct impact on the competence cascade of *S. mutans* [50]. This phenomenon was evidenced in the results of the experiments carried out by exposing the cells to low pH conditions. Similarly, oxygen is also considered a key factor which significantly alters *S. mutans* transcriptional regulation. It is found to strongly alter the competition by decreasing bacteriocins expression [51]. Carbohydrate source was as well found to have a considerable effect on the progression of the competent state of cells [50]. Hence these studies approve that the *S. mutans* has evolved from a network of regulators to integrate its cellular response to environmental change [52].

6. PHENOTYPIC DIVERSITY

As a consequence of the differences in genetic content and gene regulation, *S. mutans* strains exhibit phenotypic variability [18-20]. Around 52 different *S. mutans* genotypes have been reported in saliva and oral biofilms, with the existence of 1–5 different genotypes of *S. mutans* species in the same individual [53,54]. It has been suggested that not all strains have the same virulence capability to encourage the development of dental caries [20,22]. So, practically some of the genes related to virulence are not distributed in the same way in the strains according to their caries status [23]. Acidogenicity, aciduricity, and adhesion properties of *S. mutans* are considered as the key virulence factors [3,55,56]. The study of these virulence factors and their correlations with other species present in the biofilm is crucial in understanding the role of colonization of multiple genotypes in the same individual [53].

7. ACIDOGENECITY

The ability to produce organic acids from the carbohydrate metabolism under anaerobic conditions, with a drop in the pH value below 4.0 [3,57]. Expression of the genes related to carbohydrate absorption and metabolism seem to be responsible for the effective adaptation of *S. mutans* in the oral cavity. Carbohydrate fermentation is the key strategy for survival of *S. mutans*. They have been found to have all the genes necessary for a complete glycolytic pathway and they can produce acids even when the pH of the oral cavity is relatively low. There is sufficient evidence that *S. mutans* is one of the most acidogenic species inhabiting the dental

biofilm [58]. They can also lower the dental biofilm pH due to the intracellular polysaccharide (IPC) metabolism in the absence of fermentable carbohydrates from the diet. This was confirmed in a study by Harris et al. where they generated an IPC-deficient strain (SMS203) from the cariogenic strain UA130 and assessed its acidogenicity. It was found to be lower than the progenitor strain, thereby resulting in reduced cariogenic potential [59].

8. ACIDURICITY

It is the ability to survive rapid and extreme changes in pH and is considered to be one of the most important attributes of cariogenic bacteria. *S. mutans* can grow and carry out glycolysis at low pH values, thereby establishing a selective advantage over less aciduric species [60,61]. In order to do this, *S. mutans* are able to modify its physiology and survive in these environments due to its adaptive acid tolerance response (ATR) mechanism. This is possible due to at higher levels of expression of the proton translocator F₀F₁-ATPase pumps [60,62]. This pump works by preserving a more alkaline cytoplasmic pH compared to the extracellular environment by transporting protons out of the cell and also confers protection to acid-sensitive glycolytic enzymes [60,63,64]. Acidogenicity and aciduricity can be considered as the main factors contributing to *S. mutans* cariogenicity, as variations in these characteristics could help explain the differences in virulence among clinical isolates [20,55,56].

9. ADHESION

Colonization and survival of *S. mutans* is facilitated by its adherence to the tooth surface [60]. This adhesion, can be mediated by two mechanisms: sucrose-independent and sucrose-dependent mechanisms [55,65]. The first mechanism is not relevant for *S. mutans* virulence [56,60]. However, the sucrose-dependent mechanism is considered to be responsible for colonization of oral cavity.[55]The enzyme glycosyltransferase(GTFs), hydrolyses sucrose into glucans and fructans, The extracellular polysaccharide, glucan, not only promotes adhesion of bacteria to tooth surfaces but also promote aggregation and coaggregation of microorganisms favouring biofilm formation [55,56,65]. It has been found that *S. mutans* possesses three GTFs encoded by the *gtfB*, *gtfC*, and *gtfD* genes [55,66]. The genes encoding *gtfB* and *gtfC* are close to each other,

and are subject to the same regulatory processes. These genes are found to be expressed in response to excess glucose or sucrose [56]. The study by Yamashita et al. demonstrated that the loss of any of these genes resulted in reduction in the virulence of *S. mutans* [67].

10. CARIES AND GENOTYPIC DIVERSITY STUDIES

Pieralisi FJS et al. and Zhou Q et al. investigated the genotypic diversity of *S. mutans* in children with and without early childhood caries (ECC) and found a strong genetic diversity among *S. mutans* strains. They also stated that caries active children carried more genotypes than caries free children [68,69].

Valdez RMA et al. studied the genotypic diversity and phenotypic traits of *S. mutans* isolates and concluded that there were no differences in genotypic diversity among CF, ECC, and severe ECC children. However, the phenotypic traits such as, acidogenicity and aciduricity were found to be high in children with S-ECC [70].

Napimoga et al evaluated the relationship between clonal diversity and some virulence traits of *S. mutans* isolated from caries-free and caries-active subjects. It was found that there was larger number of genotypes of *S. mutans* with increased ability to synthesize water insoluble glucans in caries-active individuals [53].

Ravikumar et al. in 2021 reviewed seven studies related to *S. mutans* genotypic diversity and stated that the number of genotypes of *S. mutans* varied between caries active and caries-free children. They also insisted on the need for further studies to draw a definitive conclusion [10].

11. CONCLUSION

This review highlights the complexity and dynamic changes in the genetic and physical components of *S. mutans*. It is obvious that *S. mutans* has evolved numerous strategies to become well established in the oral biofilms, to antagonize the growth of commensals, to produce organic acids from dietary carbohydrates, and to propagate and metabolize under acidic conditions thereby favouring the initiation and progression of dental caries. The overall findings indicate that the acquisition of virulence genes is only a first step on the path

towards the *S. mutans* pathogenic lifestyle, it additionally requires mild genetic changes mediated by regulatory genes to adapt the expression of phenotypes with pathogenic potential according to the environmental pressures. The association between number, genotypic diversity, and caries status of an individual is still controversial. Future studies should explore the possibility of simultaneous action of different genotypes, with different phenotypic potentials, resulting in different virulence characteristics, which ultimately alters the risk of developing caries.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Fejerskov O. Changing paradigms in concepts on dental caries: Consequences for oral health care. *Caries Res.* 2004;38:182–191.
2. Takahashi N, Nyvad B. Caries ecology revisited: Microbial dynamics and the caries process. *Caries Res.* 2008;42:409–418.
3. Loesche WJ. Role of *Streptococcus mutans* in human dental decay. *Microbiol Rev.* 1986;50:353–380.
4. Tanzer JM, Livingston J, Thompson AM. The microbiology of primary dental caries in humans. *J Dent Educ.* 2001;65:1028–1037.
5. Hamada S, Slade HD. Biology, immunology, and cariogenicity of *Streptococcus mutans*. *Microbiol Rev.* 1980;44:331–384.
6. Banas et al. Glucan-binding proteins of the oral streptococci. *Crit. Rev. Oral. Biol. Med.* 2003;14(2):89–99.
7. Carlsson P, Olsson B, Bratthall D. The relationship between the bacterium *Streptococcus mutans* in the saliva and dental caries in children in Mozambique. *Arch Oral Biol.* 1985;30:265–268.
8. Matee MI, Mikx FH, de Soet JS, Maselle SY, de Graaff J, van Palenstein Helderman WH. Mutans streptococci in caries-active and caries-free infants in Tanzania. *Oral Microbiol Immunol* 1993;8:322–324.
9. Gerstein MB, et al. What is a gene, post-Encode? History and updated definition. *Genome. Res.* 2007;17(6):669–681.
10. Ravikumar D, Ramani P, Gayathri R. Genotypic diversity of *Streptococcus mutans* in children with and without early childhood caries- A systematic review. *J Oral Biol Craniofac Res.* 2021;11(2):308–312.
11. Roberts AP, Kreth J. The impact of horizontal gene transfer on the adaptive ability of the human oral microbiome. *Front Cell Infect Microbiol.* 2014;4:124.
12. Dobrindt U, Hacker J. Whole genome plasticity in pathogenic bacteria. *Curr Opin Microbiol.* 2001;4:550–7.
13. Waterhouse C, Russell RR. Dispensable genes and foreign DNA in *Streptococcus mutans*. *Microbiology.* 2006;152:1777–88.
14. Tettelin H, Riley D, Cattuto C, Medini D. Comparative genomics: The bacterial pan-genome. *Curr Opin Microbiol.* 2008;12:472–77.
15. Cornejo OE, Lefébure T, Bitar PD, Lang P, Richards VP, Eilertson K, Do T, Beighton D, Zeng L, Ahn SJ, Burne RA, Siepel A, Bustamante CD, Stanhope M. Evolutionary and population genomics of the cavity causing bacteria *Streptococcus mutans*. *Mol Biol Evol.* 2013;30:881–93.
16. Juhas M, van der Meer JR, Gaillard M, Harding RM, Hood DW, Crook DW. Genomic islands: tools of bacterial horizontal gene transfer and evolution. *FEMS Microbiol Rev.* 2009;33:376–93.
17. Koehler B, Andreen I, Jonsson B. The earlier the colonization by mutans streptococci, the higher the caries prevalence at 4 years of age. *Oral Microbiol Immunol.* 1988;3:14–17.
18. Nakano K, Nomura R, Nakagawa I, Hamada S. Demonstration of *Streptococcus mutans* with a cell wall polysaccharide specific to a new serotype,

- k, in the human oral cavity demonstr. J Clin Microbiol. 2004;42:198–202.
19. Balakrishnan M, Simmonds RS, Kilian M, Tagg JR. Different bacteriocin activities of *Streptococcus mutans* reflect distinct phylogenetic lineages. J Med Microbiol. 2002;51:941–8.
 20. Lembo FL, Longo PL, Ota-Tsuzuki C, Rodrigues CR, Mayer MP. Genotypic and phenotypic analysis of *Streptococcus mutans* from different oral cavity sites of caries-free and cariesactive children. Oral Microbiol Immunol. 2007;22:313–9.
 21. Ajdic D, McShan WM, McLaughlin RE, Savic G, Chang J, Carson MB, Primeaux C, Tian R, Kenton S, Jia H, Lin S, Qian Y, Li S, Zhu H, Najjar F, Lai H, White J, Roe BA, Ferretti J. Genome sequence of *Streptococcus mutans* UA159, a cariogenic dental pathogen. Proc Natl Acad Sci U S A. 2002;99:14434–9.
 22. Fitzgerald DB, Fitzgerald RJ, Adams BO, Morhart R. Prevalence, distribution of serotypes, and cariogenic potential in hamsters of mutans streptococci from elderly individuals. Infect Immun. 1983;41:691–7.
 23. Argimón S, Caufield PW. Distribution of putative virulence genes in *Streptococcus mutans* strains does not correlate with caries experience. J Clin Microbiol. 2011;49:984–92.
 24. Póntigo F, Moraga M, Flores SV. Molecular phylogeny and a taxonomic proposal for the genus *Streptococcus*. Genet Mol Res. 2015;14:10905–18.
 25. Lal D, Verma M, Lal R. Exploring internal features of 16S rRNA gene for identification of clinically relevant species of the genus *Streptococcus*. Ann Clin Microbiol Antimicrob. 2011;10:28.
 26. Whiley RA, Beighton D. Current classification of the oral streptococci. Oral Microbiol Immunol. 1998;13:195–216.
 27. Shibata Y, Ozaki K, Seki M, Kawato T, Tanaka H, Nakano Y, Yamashita Y. Analysis of loci required for determination of serotype antigenicity in *Streptococcus mutans* and its clinical utilization. J Clin Microbiol. 2003;41:4107–12.
 28. Nakano K, Nomura R, Matsumoto M, Ooshima T. Roles of oral bacteria in cardiovascular diseases— from molecular mechanisms to clinical cases: cell-surface structures of novel serotype k *Streptococcus mutans* strains and their correlation to virulence. J Pharmacol Sci. 2010;113:120–5.
 29. Nakano K, Nomura R, Shimizu N, Nakagawa I, Hamada S, Ooshima T. Development of a PCR method for rapid identification of new *Streptococcus mutans* serotype k strains. J Clin Microbiol. 2004;42:4925–30.
 30. Lapirattanakul J, Nakano K, Nomura R, Nemoto H, Kojima A, Senawongse P, Srisatjaluk R, Ooshima T. Detection of serotype k *Streptococcus mutans* in Thai subjects. Oral Microbiol Immunol. 2009;24:431–3.
 31. Song L, Wang W, Conrads G, Rheinberg A, Sztajer H, Reck M, Wagner-Döbler I, Zeng A. Genetic variability of mutans streptococci revealed by wide whole-genome sequencing. BMC Genomics. 2013;14:430.
 32. Shields RC, Zeng L, Culp DJ, Burne RA. Genomewide identification of essential genes and fitness determinants of *Streptococcus mutans* UA159. MSphere. 2018;3:e00031-18.
 33. Lorenz MG, Wackernagel W. Bacterial gene transfer by natural genetic transformation in the environment. Microbiol Rev. 1994;58:563–602.
 34. Maruyama F, Kobata M, Kurokawa K, Nishida K, Sakurai A, Nakano K, Nakagawa I. Comparative genomic analyses of *Streptococcus mutans* provide insights into chromosomal shuffling and species-specific content. BMC Genomics. 2009;10:358.
 35. Ferretti JJ, Ajdic D, McShan WM. Comparative genomics of streptococcal species. Indian J Med Res. 2004;119 Suppl:1–6.
 36. Periwal V, Scaria V. Insights into structural variations and genome rearrangements in prokaryotic genomes. Bioinformatics. 2015;31:1–9.
 37. Hastings PJ, Lupski JR, Rosenberg SM, Ira G. Mechanisms of change in gene copy number. Nat Rev Genet. 2009;10:551–64.
 38. Meng P, Lu C, Zhang Q, Lin J, Chen F. Exploring the genomic diversity and cariogenic differences of *Streptococcus mutans* strains through pan-genome and comparative genome analysis. Curr Microbiol. 2017;74:1200–9.
 39. Hoshino T, Fujiwara T, Kawabata S. Evolution of cariogenic character in *Streptococcus mutans*: Horizontal

- transmission of glycosyl hydrolase family 70 genes. *Sci Rep.* 2012;2:518.
40. Argimón S, Alekseyenko AV, DeSalle R, Caufield PW. Phylogenetic analysis of glucosyltransferases and implications for the coevolution of mutans streptococci with their mammalian hosts. *PLoS One.* 2013;8:e56305.
 41. Rouli L, Merhej V, Fournier PE, Raoult D. The bacterial pangenome as a new tool for analysing pathogenic bacteria. *New Microbes New Infect.* 2015;7:72–85.
 42. Zhu W, Liu S, Liu J, Zhou Y, Lin H. High-throughput sequencing identification and characterization of potentially adhesion-related small RNAs in *Streptococcus mutans*. *J Med Microbiol.* 2018;67:641–51.
 43. Shields RC, O'Brien G, Maricic N, Kesterson A, Grace M, Hagen SJ, et al. Genome-wide screens reveal new gene products that influence genetic competence in *Streptococcus mutans*. *J Bacteriol.* 2017;200:e00508-17.
 44. Mattos-Graner RO, Duncan MJ. Two-component signal transduction systems in oral bacteria. *J Ora Microbiol.* 2017;9:1400858.
 45. Zheng L, Chen Z, Itzek A, Herzberg M, Kreth J. CcpA regulates biofilm formation and competence in *Streptococcus gordonii*. *Mol Oral Microbiol.* 2012;27:83–94.
 46. Mahillon J, Chandler M. Insertion sequences. *Microbiol Mol Biol Rev.* 1998;62:725–74.
 47. Palmer SR, Miller JH, Abranches J, Zeng L, Lefebure T, Richards VP, Lemos AJ, Stanhope MJ, Burne R. Phenotypic heterogeneity of genomically-diverse isolates of *Streptococcus mutans*. *PLoS One.* 2013;8(4):e61358.
 48. Merritt J, Qi F. The mutacins of *Streptococcus mutans*: regulation and ecology. *Mol Oral Microbiol.* 2012;27:57–69.
 49. Federle MJ, Morrison D. One if by land, two if by sea: signaling to the ranks with CSP and XIP. *Mol Microbiol.* 2012;86:241–5.
 50. Moye Z., Son M, Rosa-Alberty A E, Zeng L, Ahn SJ, Hagen SJ, Burne RA. Effects of carbohydrate source on genetic competence in *Streptococcus mutans*. *Appl Environ Microbiol.* 2016;82:4821–34.
 51. Ahn SJ, Wen ZT, Burne RA. Effects of oxygen on virulence traits of *Streptococcus mutans*. *J Bacteriol.* 2007;189:8519–27.
 52. Smith EG, Spatafora GA. Gene regulation in *S. mutans*: Complex control in a complex environment. *J Dent Res.* 2012;91:133–41.
 53. Napimoga MH, Höfling JF, Klein MI, Kamiya RU, Gonçalves RB. Transmission, diversity and virulence factors of *Streptococcus mutans* genotypes. *J Oral Sci.* 2005;47:59–64.
 54. Tabchoury CP, Sousa MC, Arthur RA, Mattos-Graner RO, Del Bel Cury AA, Cury J. Evaluation of genotypic diversity of *Streptococcus mutans* using distinct arbitrary primers. *J Appl Oral Sci.* 2008;16:403–7.
 55. Banas JA. Virulence properties of *Streptococcus mutans*. *Front Biosci.* 2004;9:1267–77.
 56. Krzyściak W, Jurczak A, Kościelniak D, Bystrowska B, Skalniak A. The virulence of *Streptococcus mutans* and the ability to form biofilms. *Eur J Clin Microbiol Infect Dis.* 2014;33:499–515.
 57. Nishimura J, Saito T, Yoneyama H, Lan Bai L, Okumura K, Isogai E. Biofilm formation by *Streptococcus mutans* and related bacteria. *Adv Microbiol.* 2012;2:208–15.
 58. De Soet JJ, Nyvad B, Kilian M. Strain-related acid production by oral streptococci. *Caries Res.* 2000;34:486–90.
 59. Harris GS, Michalek SM, Curtiss R. Cloning of a locus involved in *Streptococcus mutans* intracellular polysaccharide accumulation and virulence testing of an intracellular polysaccharide deficient mutant. *Infect Immun.* 1992;60:3175–85.
 60. Lemos, J.A., Abranches, J., Burne RA. Responses of cariogenic streptococci to environmental stresses. *Curr Issues Mol Biol.* 2005;7:95–107.
 61. Matsui R, Cvitkovitch D. Acid tolerance mechanisms utilized by *Streptococcus mutans*. *Future Microbiol.* 2010;5:403–17.
 62. Nascimento MM, Lemos JA, Abranches J, Gonçalves RB, Burne R. Adaptive acid tolerance response of *Streptococcus sobrinus* adaptive acid tolerance response of *Streptococcus sobrinus*. *J Bacteriol.* 2004;186:6383–90.
 63. Van de Guchte M, Serror P, Chervaux C, Smokvina T, Ehrlich SD, Maguin E. Stress

- responses in lactic acid bacteria. Antonie Van Leeuwenhoek. 2002;82:187–216.
64. Baker JL, Faustoferrri RC, Quivey RG. Acid-adaptive mechanisms of *Streptococcus mutans*-the more we know, the more we don't. *Mol Oral Microbiol*. 2017;32:107–17.
 65. Bowen WH, Koo H. Biology of *Streptococcus mutans*-derived glucosyltransferases: role in extracellular matrix formation of cariogenic biofilms. *Caries Res*. 2011;45:69–86.
 66. Munro CL, Michalek SM, Macrina F. Sucrose-derived exopolymers have site-dependent roles in *Streptococcus mutans*-promoted dental decay. *FEMS Microbiol Lett*. 1995;128:327–32.
 67. Yamashita Y, Bowen WH, Burne RA, Kuramitsu H. Role of the *Streptococcus mutans* *gtf* genes in caries induction in the specific-pathogen-free rat model. *Infect Immun*. 1993;61:3811–7.
 68. Peralisi FJ, Rodrigues MR, Segura VG, Maciel SM, Ferreira FB, Garcia JE, Poli-Frederico RC. Genotypic Diversity of *Streptococcus mutans* in Caries-Free and Caries-Active Preschool Children. *Int J Dent*. 2010;2010:824976.
 69. Zhou Q, Qin X, Qin M, Ge L. Genotypic diversity of *Streptococcus mutans* and *Streptococcus sobrinus* in 3-4-year-old children with severe caries or without caries. *Int J Paediatr Dent*. 2011;21(6): 422-31.
 70. Valdez RMA, Duque C, Caiiffa KS, Dos Santos VR, Loesch MLA, Colombo NH, Arthur RA, Negrini TC, Boriollo MFG, Delbem ACB. Genotypic diversity and phenotypic traits of *Streptococcus mutans* isolates and their relation to severity of early childhood caries. *BMC Oral Health*. 2017;17(1): 115.

© 2021 Supraja et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/82438>