



Oxidative Effects in Streptozotocin-induced Male and Female Mice: The Effect of Garlic Oil and Melatonin

Abeer Al-Rashedi ^{a#}, Banan Al-Johari ^{a#}, Rahaf Al-Abbasi ^{a#}, Raghda Sindi ^{a#},
Rawan Bin Hameed ^{a#}, Sulafah Al-Saadi ^{a#}, Sara Al-Awam ^{a#},
Mawada Bukhari ^{a#}, Rahaf Al-Talhi ^{a#} and Yosra Al-Hindi ^{b*†}

^a Faculty of Pharmacy, Umm Al-Qura University, KSA, Saudi Arabia.
^b Clinical Pharmacy, Faculty of Pharmacy, Umm Al-Qura, Saudi Arabia.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i58A34124

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/79896>

Original Research Article

Received 10 October 2021
Accepted 14 December 2021
Published 15 December 2021

ABSTRACT

Background: Recent studies have revealed that a hyperglycemia-induced overproduction of superoxide can be the first event in the activation of all pathways involved in the pathogenesis of complications of diabetes. Supplementation of garlic was found to decrease diabetes-induced oxidative stress complications. Studies shown also that melatonin attenuates diabetes-induced oxidative stress in diabetic induced rabbits and rats.

Objective: In this present study, oxidative stress in diabetic model and the effect of garlic oil or melatonin treatment were examined in both genders' male and females' mice.

Methods: 96 mice were randomly divided into 5 groups including control (C), diabetic (D), melatonin 10 mg/kg (D+M), garlic extract 100 mg/kg (D+G) and combined melatonin and garlic (D+M+G). All treatments were given orally daily for 16 weeks after induction of hyperglycemia by streptozotocin (STZ). Fasting blood glucose and antioxidant levels were estimated.

Results: Streptozotocin induced diabetic mice, showed a significant increase of plasma glucose, lipid peroxide and uric acid. Accordingly, significant decreases in the levels of antioxidants

Pharm D undergraduate;

† Assistant Professor;

*Corresponding author: E-mail: yzhindi@uqu.edu.sa;

ceruloplasmin were found in the plasma of diabetic mice. Treatment of diabetic mice with garlic oil or melatonin for 16 weeks significantly increased plasma levels of ceruloplasmin activities. Lipid peroxides, uric acid, blood glucose was decreased significantly after treatment with garlic oil or melatonin.

Conclusion: The results suggest that garlic oil or melatonin may effectively normalize the impaired antioxidants status in streptozotocin induced diabetes in both males and females mice.

Keywords: Diabetes; Garlic; Melatonin; STZ; mice; antioxidants.

1. INTRODUCTION

Nowadays, researches are focused on the relation of antioxidants activity and diabetes [1]. The oxygen species like hydrogen peroxide, superoxide anions, singlet oxygen and hydroxyl radicals could be developed through ionizing radiation and aerobic metabolism of either endogenous or exogenous materials [2]. Evidence have shown that diabetes has been related to an increased free radical production [3]. The mechanisms of this state could be contributed to the free radical development in diabetes that can increase the non-enzymatic and auto-oxidative glycosylation, and moreover metabolic stress due to differences in the metabolism of energy, inflammatory processes, and antioxidant activation [4].

A lot of focus has been developed in using herbal medicine and other natural products in treating diseases that have a state of oxidative stress like diabetes and metabolic syndrome [5]. Garlic is one of the elements that were used as herbal therapy for long time and have been seen to help in eliminating cardiovascular risks and diabetes [6]. Moreover, melatonin was found to have a great antioxidant ability towards diseases [7].

To our knowledge, no previous researches if any used the combination of garlic and melatonin in diabetic subjects to measure antioxidants activity. Therefore, due to the frequent use of antidiabetic drugs with their associated side effects, we hypothesized that combination of low doses of garlic and melatonin might have beneficial effects on glycemic and antioxidants in diabetes mellitus. Also, this combination can be used with traditional antidiabetic drugs to reduce their doses and potential side effects. The aim of this study was to investigate the effects of low doses of melatonin and garlic extract both individually and combined on blood glucose levels as well as antioxidants in streptozotocin induced diabetic male and female mice.

2. MATERIALS AND METHODS

2.1 Reagents

Reagents and chemicals including were purchased from Sigma-Aldrich (MO, USA), unless otherwise stated.

2.2 Preparation of Garlic Extract

Black garlic (BG) was supplied from Ueising-Nongsan (Korea), and extracted by heating with water twice under reflux at 80°C (yield, 12.8%). The resulting solution was freeze-dried after evaporation and kept at 4°C until use. The BG was fermented with edible *Saccharomyces cerevisiae* (KCTC 7910) by 2-stage cultivation. At the first stage cultivation, the microorganism was cultivated in a medium containing 3% (wt/vol) malt extract for 36 hours at 28°C to enhance cell growth. The cell mass was obtained by centrifugation and recultivated in a medium containing 5% (wt/vol) of the BG extracts under the same conditions to increase the concentration of physiologically active substances, such as polyphenol and allicystein, which play a key role in antioxidation. After cultivation, the culture solutions were extracted by heating after filtration to remove the cells. The solution was then freeze-dried after evaporation and kept at 4°C until needed. Then it was dissolved directly in distilled water and administered orally at a dose of 100 mg/kg, once per day for 16 weeks after induction of diabetes [8]. The S-allyl cysteine sulfoxide is responsible for anti-peroxide effect of garlic.

2.3 Melatonin

Melatonin from (Sigma-Aldrich Co., St. Louis, MO) 10 mg/kg daily dissolved in 0.04% ethanol added to the drinking water at night [9].

2.4 Animals

All procedures concerning animal care and treatment were approved by Umm Al-Qura

university's Biomedical and research Ethics Committee (HAPO-02-K-012-2021-10-788). For this study total number of 96 males and females of C57BL/6J (B6) strain mice were purchased from Harlan (Charles River Laboratories, Wilmington, Massachusetts, USA). Mice were maintained in a temperature-controlled room ($23 \pm 1^\circ\text{C}$) under a 12-h to 12-h light to dark cycle. Mice were individually housed in standard cages with *ad libitum* water and standard chow (CRM pellets, SDS diets, U.K.).

Measurements then started at the age of 10 weeks with ranging body weight of 25.7g to 30.33g and were taken over a period of 16 weeks. We choose this period of time because diabetes is a chronic disease and we wanted to explore this at acute and long-term use of these herbal medicines. After 16 weeks mice were fasted overnight and euthanized by CO₂ and blood samples were taken by cardiac puncture.

2.5 Induction and Assessment of Diabetes

A method of inducing type 2 DM with hyperglycemia and relatively low insulin levels can be produced STZ. Mice were intraperitoneally (i.p.) injected with STZ (50 mg/kg of BW) in 0.1 M citrate buffer (pH 4.2) on two consecutive days. NA (120 mg/kg of BW) in saline was i.p. injected 30 min before the STZ injection on the first day after overnight fasting. Seven days after the second i.p. injection, mice that exhibited an 8-h fasting blood glucose (FBG) level of 200 mg/dL were recognized as being hyperglycemic. The others that exhibited an FBG level of <200 mg/dL were injected with STZ and

monitored until the FBG level reached 200 mg/dL. FBG was monitored with a glucometer (Dragon Pharmaceutical Co, New Taipei, Taiwan).

2.6 Experimental Design

Mice were randomly classified into 4 groups (n=16 each, 8 Males and 8 Females); one group served as a control non-diabetic group and 3 groups were injected with STZ to induce diabetes, then they received their respective drug treatments daily for 16 consecutive weeks according to the following design (Fig. 1):

- Group 1(C); control non-diabetic mice treated with saline orally and citrate buffer intraperitoneally.
- Group 2 (D); nontreated induced diabetic mice.
- Group 3 (D+M); induced diabetic mice treated with melatonin [9] (10 mg/kg/day in drinking water)
- Group 4 (D+G); induced diabetic mice treated with prepared garlic extract [8] (100 mg/kg/day in drinking water)

The doses of melatonin and garlic were chosen based on previous experimental studies that examined wide range of doses in diabetes. The lowest doses were selected for both melatonin and garlic due to the potential side effects of large doses.

At 2nd and 16th weeks, the levels of FBG were examined. After 16 weeks antioxidants were examined.




Fig. 1. Experimental design of the study

2.7 Assessment

2.7.1 Fasting blood glucose measurements

At week 2 and 16 all mice were fasted overnight and blood were taken from their tails in order to measure their FBG and insulin levels per the manufacturer's instructions. Fasting blood glucose was measured with a One Touch II glucose meter (Lifescan, Inc., Johnson & Johnson, Milpitas, CA).

2.7.2 Antioxidant's measurements

Lipid peroxide levels were measured in plasma, hemolysate and tissue homogenates as thiobarbituric acid reactivity (TBARS) [10]. The product of the reaction between malondialdehyde and thiobarbituric acid was measured as described by Thayer. Ceruloplasmin activity was determined using a para-phenylenediamine dihydrochloride method [10]. Uric acid was determined by enzymatic colorimetric method using commercial kit (Biocon, BurbachyGermany) [10].

2.7.3 Statistical analysis

All results are expressed as group means \pm SEM. Results were analyzed by one-way analysis of variance, followed by Tukey's post-

hoc test to assess significance, using a criterion of P value of less than 0.05. The statistical analysis was carried out using GraphPad Prism version 5 (GraphPad Software Inc., California, USA).


3. RESULTS

3.1 Effects of Melatonin and Garlic on Fasting Blood Glucose in Diabetic Mice

Fig. 2 describes the effect of garlic, melatonin and their combination on fasting blood glucose on both male and female mice. It is significant that after 16 week there were a vast improvement in all treatments in decreasing blood glucose levels in diabetic mice ($P < 0.005$).

3.2 Effects of Melatonin and Garlic on Antioxidant levels in Diabetic Mice

Fig. 3 describes the effect of garlic, melatonin and their combination on antioxidants levels on both male and female mice. It is significant that after 16 weeks of treatments there was an increased plasma level of ceruloplasmin activities. Lipid peroxides and uric acid were decreased significantly after treatment with garlic oil or melatonin or their combinations ($P < 0.005$).







Fig. 2. Effects of Melatonin and Garlic on Fasting Blood Glucose in Diabetic Mice; A) Plasma Glucose measurements in week 2 of the study. B) Plasma Glucose measurements in week 16 of the study C; control, D; diabetic induced mice, D+M; diabetic mice treated with melatonin, D+G; diabetic mice treated with garlic, D+M+G; diabetic mice treated with combination of melatonin and garlic

* $P < 0.05$ compared to control mice, ** $P < 0.05$ compared to diabetic mice






Fig. 3. Effects of Melatonin and Garlic on antioxidant levels in Diabetic Mice; A) Levels of lipids peroxidase; B) Ceruloplasmin and C) Uric acids after 16 weeks of treatments

* $P < 0.05$ compared to control mice, ** $P < 0.05$ compared to diabetic mice, + $P < 0.001$ compared to D+M+G

4. DISCUSSION

Diabetes can cause multiple changes systemically and also upon the cellular levels [10,11]. In our present study, diabetic mice both genders induced by streptozotocin showed the expected raise in their plasma glucose levels and the decrease in antioxidant levels. However, treatment of diabetic mice with garlic oil or melatonin for 16 weeks significantly increased plasma levels of ceruloplasmin activities. Lipid peroxides, uric acid, blood glucose was decreased significantly after treatment with garlic oil or melatonin. Our results were in agreement of other researchers whom used garlic or melatonin alone, they showed a lower levels of blood glucoses and other parameters as cholesterol and lipid profile [11,12,13,14]. Researches tried to explain the possible mechanism of this blood glucose lowering, for garlic they approved multiple pathways of anti-inflammatory and antihyperlipidemic and antioxidant effects. On the other hand, for the melatonin they believed it works on brain site and can achieves this action through direct detoxification of reactive oxygen and reactive nitrogen species and indirectly by stimulating antioxidant enzymes at the same time inhibiting the activity of enzymes as prooxidant's. Moreover, melatonin shown to chelates transition metals, that have an involvement in the Fenton/Haber–Weiss reactions; therefore, melatonin can decrease toxic hydroxyl radical that results in the inhibition of oxidative stress. However, more investigation for a clearer explanation is in need [15]. Evidence showed

also that when given melatonin to rats taking cytotoxic drugs that there state of antioxidants were improved [16,17,18,19].

There is evidence from a study done on antiperoxide effect of garlic oil on various tissues and blood cell counts of arsenic exposed albino mice found that there was a significant reduction in the arsenic levels. That indicates garlic pearls can be administered to persons who are working in places of coal burning, mining and metal ore smelting, as the diallyl disulphide (DADS) that is the active principle of garlic has a possible protective role on LPO status of many tissues and blood cell counts of arsenic exposed mice [20].

From the benefits shown in our study of using the combination of garlic and melatonin together in diabetic subjects, we recommend starting a trail with diabetic patients using the combination of melatonin and garlic to investigate in depth their beneficial effect.

5. CONCLUSION

Our results from this present study suggest that garlic oil or melatonin may effectively normalize the impaired antioxidants status in streptozotocin induced diabetes in both males and females mice. The effects of these antioxidants of both agents may be useful in delaying the complications of diabetes such as retinopathy, nephropathy and neuropathy due to imbalance between free radicals and antioxidant systems.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

CONSENT

It is not applicable.

ETHICAL APPROVAL

All procedures concerning animal care and treatment were approved by Umm Al-Qura university's Biomedical and research Ethics Committee (HAPO-02-K-012-2021-10-788).

The study highlights the efficacy of " Herbal " which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

DATA AVAILABILITY

The datasets generated and/or analysed during the current study are available from the corresponding author on reasonable request.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Malode LL, Manwar JV, Panchale WA, Bartere SA, Bakal RL. Potential of medicinal plants in management of diabetes: An updates. *GSC Advanced Research and Reviews*. 2021;8(1):149-59.
2. Costa TJ, Barros PR, Arce C, Santos JD, da Silva-Neto J, Egea G, Dantas AP, Tostes RC, Jimenez-Altayo F. The homeostatic role of hydrogen peroxide, superoxide anion and nitric oxide in the vasculature. *Free Radical Biology and Medicine*. 2021;162:615-35.
3. Davison GW, Irwin RE, Walsh CP. The metabolic-epigenetic nexus in type 2 diabetes mellitus. *Free Radical Biology and Medicine*; 2021.
4. Carvalho MG, Silva KM, Aristizabal VH, Ortiz PE, Paranzini CS, Melchert A, Amaro JL, Souza FF. Effects of Obesity and Diabetes on Sperm Cell Proteomics in Rats. *Journal of Proteome Research*. 2021;20(5):2628-42.
5. Adhikari B, Marasini BP, Rayamajhee B, Bhattarai BR, Lamichhane G, Khadayat K, Adhikari A, Khanal S, Parajuli N. Potential roles of medicinal plants for the treatment of viral diseases focusing on COVID-19: A review. *Phytotherapy Research*. 2021;35(3):1298-312.
6. Reiter RJ, Mayo JC, Tan DX, Sainz RM, Alatorre-Jimenez M, Qin L. Melatonin as an antioxidant: under promises but over delivers. *Journal of pineal research*. 2016;61(3):253-78.
7. Abdollahzade N, Majidinia M, Babri S. Melatonin: A pleiotropic hormone as a novel potent therapeutic candidate in arsenic toxicity. *Molecular Biology Reports*. 2021 Sep;48(9):6603-18.
8. Thomson M, Al-Qattan KK, Divya JS, Ali M. Anti-diabetic and anti-oxidant potential of aged garlic extract (AGE) in streptozotocin-induced diabetic rats. *BMC Complementary and Alternative Medicine*. 2016;16(1):1-9.
9. Korac B, Kalezic A, Pekovic-Vaughan V, Korac A, Jankovic A. Redox changes in obesity, metabolic syndrome, and diabetes. *Redox Biology*. 2021;101887.
10. Anwar MM, Meki AR. Oxidative stress in streptozotocin-induced diabetic rats: effects of garlic oil and melatonin. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*. 2003;135(4):539-47.
11. Lade PN, Naidu K, Yenarkar S. Benefits of Using Black Garlic and its Role In Treating Liver Diseases Along With Boosting Immunity.
12. Faran SA, Asghar S, Khalid SH, Khan IU, Asif M, Khalid I, Farooq Gohar U, Hussain T. Hepatoprotective and renoprotective properties of lovastatin-loaded ginger and garlic oil nanoemulsomes: insights into serum biological parameters. *Medicina*. 2019;55(9):579.
13. Lo CC, Lin SH, Chang JS, Chien YW. Effects of melatonin on glucose homeostasis, antioxidant ability, and

- adipokine secretion in ICR mice with NA/STZ-induced hyperglycemia. *Nutrients*. 2017;9(11):1187.
14. Mumtaz S, Ali S, Khan R, Shakir HA, Tahir HM, Mumtaz S, Andleeb S. Therapeutic role of garlic and vitamins C and E against toxicity induced by lead on various organs. *Environmental Science and Pollution Research*. 2020;27(9):8953-64.
 15. Asgary S, Rastqar A, Keshvari M. Functional food and cardiovascular disease prevention and treatment: a review. *Journal of the American College of Nutrition*. 2018;37(5):429-55
 16. Pérez-López FR, Chedraui P, Haya J, Cuadros JL. Effects of the Mediterranean diet on longevity and age-related morbid conditions. *Maturitas*. 2009;64(2):67-79.
 17. Nishida S, Segawa T, Murai I, Nakagawa S. Long-term melatonin administration reduces hyperinsulinemia and improves the altered fatty-acid compositions in type 2 diabetic rats via the restoration of Δ -5 desaturase activity. *Journal of pineal research*. 2002;32(1):26-33.
 18. Anisimov VN, Popovich IG, Zabezhinski MA, Anisimov SV, Vesnushkin GM, Vinogradova IA. Melatonin as antioxidant, geroprotector and anticarcinogen. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*. 2006;1757(5-6):573-89.
 19. Altun A, Ugur-Altun B. Melatonin: therapeutic and clinical utilization. *International journal of clinical practice*. 2007;61(5):835-45.
 20. R. Latha, L. Venkatakrisnan, V. Aruna, G. Lavanya. The effect of garlic oil on lipid peroxidation and blood cell counts of arsenic exposed albino mice. *Journal of Natural Remedies* 2006;6(1): 19-25.

© 2021 Al-Rashedi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/79896>