



Lipid Peroxidation and Glutathione Level Following Scopolamine-Induced Cognitive Dysfunction in Rats: Potentials of *Telfairia occidentalis* Seeds and *Talinium triangulare* Leaves Aqueous Extracts

Samson O. Paulinus ^a, Eru M. Eru ^{b*},
Gabriel U. Udo-Affah ^b, Michael E. Oku ^b, Ifiok Bassey ^b,
Kelechi C. Uruakpa ^b, Nsikak M. Umoh ^b, Michael Ajen ^c,
Sadeyeng E. Anani ^b and Anozeng O. Igiri ^b

^a Department of Radiography & Radiological Science, University of Calabar, Calabar, Nigeria.

^b Department of Anatomy, University of Calabar, Calabar, Nigeria.

^c Department of Biochemistry, University of Calabar, Calabar, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121691>

Original Research Article

Received: 23/06/2024

Accepted: 26/08/2024

Published: 30/08/2024

*Corresponding author: E-mail: erumba@unical.edu.ng;

Cite as: Paulinus, Samson O., Eru M. Eru, Gabriel U. Udo-Affah, Michael E. Oku, Ifiok Bassey, Kelechi C. Uruakpa, Nsikak M. Umoh, Michael Ajen, Sadeyeng E. Anani, and Anozeng O. Igiri. 2024. "Lipid Peroxidation and Glutathione Level Following Scopolamine-Induced Cognitive Dysfunction in Rats: Potentials of *Telfairia Occidentalis* Seeds and *Talinium Triangulare* Leaves Aqueous Extracts". *Asian Journal of Research and Reports in Neurology* 7 (1):98-106. <https://www.journalajorrin.com/index.php/AJORRIN/article/view/107>.

ABSTRACT

Background: Oxidative stress augmented with progressive age, causes changes in mitochondrial DNA, mitochondrial disruption and more oxidative trouble. This process is facilitated in Alzheimer's disease by the pathology of $\alpha\beta$ amyloid and activated microglia.

Objectives: The present study compared the potentials of aqueous extract of *Telfairia occidentalis* (TO) seeds and *Talinum triangulare* (TT) on the malondialdehyde (MDA) and glutathione (GSH) levels following scopolamine hydrobromide (SHB)-induced Alzheimer's type cognitive dysfunction in rats.

Methods: Forty-two Wistar rats were arrayed into seven arrays (I-VII). Alzheimer's type cognitive dysfunction was induced in arrays II-VII by administering intraperitoneally (IP) 1 mg/kg body weight (BW) of SHB for seven days before aqueous extracts of TO (850 mg/kg and 1750 mg/kg), TT (850 mg/kg and 1750 mg/kg) and donepezil (1 mg/kg) administrations for 14 days. The rats' blood serums were taken and analyzed.

Result: The MDA estimation in group II (672 ± 39.65), VI (707.67 ± 24.99) and VII (671.64 ± 32.07) increased significantly compared with others ($P<0.05$). Glutathione level was significantly increased in arrays I (0.44 ± 0.05), IV (0.41 ± 0.08), V (0.46 ± 0.07) and VII (0.44 ± 0.05) compared to others.

Conclusion: Aqueous extract of TO seeds reduced MDA levels and both extracts increased glutathione levels in Alzheimer's type cognitive dysfunction rats; though the effect of TT was dose-dependent.

Keywords: Alzheimer's disease; glutathione; lipid peroxidase; oxidative stress; wistar rats.

1. INTRODUCTION

Lipid peroxidation is an act through which oxidants such as free radicals or non-radical species fight lipids with carbon-carbon double bonds, such as polyunsaturated fatty acids [1]. At sub-toxic conditions (low lipid peroxidation), the cells actuate their maintenance and survival through antioxidant defense mechanisms or signaling pathways activation that up-regulate antioxidant proteins resulting in an adaptive stress response. At medium or elevated lipid α peroxidation states (toxic condition), the extent of oxidative stress overpowers repair capacity and causes cells' apoptosis or necrosis; both processes eventually lead to molecular cell destruction which may hasten the development of different disease states and accelerate ageing [2].

On the other hand, Glutathione directly removes different oxidants such as superoxide anion, hydroxyl radical, nitric oxide and carbon radicals. Glutathione catalytically neutralizes hydroperoxides, peroxy nitrates and lipid peroxides [3]. In health, accumulation of glutathione disulphide due to oxidative stress is directly poisonous to cells, causing programmed cell death by activation of the stress activated protein kinase/mitogen-activated protein kinase (SAPK/MAPK) pathway [4]. Glutathione depletion triggers apoptosis [3]. The disease associated with glutathione depletion includes

neurodegenerative diseases such as Alzheimer's (AD), Parkinson's and others [5].

Oxidative stress augmented with progressive age, causes changes in mitochondrial DNA, mitochondrial disruption and more oxidative trouble. This process is facilitated in AD by the act of $\alpha\beta$ and actuated microglia [6]. Oxidative trouble underlies the gradual neurodegenerative characteristics of AD [7]. Reports have it that oxidative stress elevates the expression and turns on β and γ secretase and inhibits the actions of α -secretase [8]. High formation and absence of A β peptides removal can sediment $\alpha\beta$ which energizes different cell signaling networks, hence, causing degradation of synapses, loss of neurons and reduced cognitive function [9].

In vitro experiments revealed $\alpha\beta$ elevated H_2O_2 and lipid peroxidase statuses. Age-associated $\alpha\beta$ sedimentation elevates hydrogen peroxide, nitric oxide formation and oxidative mutation of proteins and lipids deducing that $\alpha\beta$ causes oxidative stress [10,11]. In hippocampal neuron cultures, the cause of oxidants by A β needs the stimulation of N-methyl diaspertate (NMDA) receptor and quick elevated neuronal Ca^{2+} status [12]. The externally generated antioxidants are mainly from food and herbs [13]. These antioxidants abet to counteract the surplus enlarged radicals, guard the cell against envenom effects and contribute to ailment aversion [14] However, comparing the effects of

aqueous extracts of *Telfairia occidentalis* (TO) seeds and *Talinum triangulare* (TT) leaves on the lipids and glutathione levels using Alzheimer's type cognitive dysfunction rats may provide in-depth knowledge to their physiology, thus, the necessity for this research.

2. MATERIALS AND METHODS

Experimental animals: With ethical approval number: FAREC-FBMS 042ANA3719, forty-two adult Wistar rats with both sexes and weight ranging from 180-200g were bought from the animal farm and kept in the Departmental animal room for two weeks. Before the experiment, the experimental animals were kept for acclimatization under standard conditions of temperature (27°C – 30°C), given rat chow and water *ad libitum*. After two weeks of adjustment, the rats were arbitrarily arrayed into seven arrays; each having six rats designated I-VII.

Plant extract preparation: Fresh TO seeds and TT leaves were bought from the market in Cross River State, Nigeria. These seeds and leaves were registered with voucher numbers: HERB/BOT/UCC/322 and HERB/BOT/UCC/120 in Botany Department, University of Calabar, Calabar. The TO seeds were exposed from the husks, washed alongside with the TT leaves, reduced into tiny parts and dehydrated (air dried) in the laboratory. The dehydrated specimens were pulverized into powdered form (1600g) with a blender (model number Bravo3JARS Mixer grinder) and then soaked in 1000 mL of distilled water for 24 hours. The admixture was purified with Whatman No.1 filter paper and chess cloth. The products were collected and made potent to a goeey remnant at 40-50° and kept in a cool dry place for later use.

Alzheimer's type cognitive dysfunction induction: 1.0 mg/kg bw of SHB (bought from Bristol Scientific Company, Bristol Road, Apapa, Lagos- Nigeria) was intraperitoneally (IP) injected to adult Wistar rats in arrays II-VII for seven days to establish Alzheimer's type cognitive dysfunction.

LD₅₀ determination: Using Lorke's method [15], LD₅₀ of aqueous extracts of TO seeds and TT leaves were both established to be >7000 mg/kg and doses were determined using 12.5% and 25% of the established LD₅₀.

Plants extract and donepezil administration: Array I served as the negative control and received animal feed and water *ad libitum*; array

II served as the positive control and received 1.0 mg/kg body weight of SHB only; array III received 1.0 mg/kg body weight of SHB and 1.0 mg/kg body weight of Donepezil (bought from Bez Pharmacy, Etta Agbo Calabar-Nigeria); array IV received 1.0 mg/kg body weight of SHB and 875 mg/kg body weight of aqueous TO seeds; array V received 1.0 mg/kg body weight of SHB and 1750 mg/kg body weight of aqueous TO seeds; array VI received 1.0 mg/kg body weight of SHB and 875 mg/kg body weight of aqueous TT leaves while array VII received 1.0 mg/kg body weight of SHB and 1750 mg/kg body weight of aqueous TT leaves. These extracts and drugs were administered for fourteen days.

Determination of MDA and glutathione: 24 hours after the last extracts and drug administration, the experimental animals were sacrificed and the blood serums were taken through cardiac puncture for analysis. Serum levels of Lipid peroxidation and glutathione peroxidase were used as markers for oxidative stress. Lipid peroxidation was estimated by measuring 0.25ml of serum and 1.25ml of 10% trichloroacetic acid and added to a clean centrifuge tube and allowed for 10 minutes. 1.25mls of 0.05 M H₂SO₄ and 1.5ml of 0.67 TBA (thiobarbituric acid) were placed in a boiling water tap and 2ml of butanol was added. Thiobarbituric acid reactive material was extracted, and absorbance was read at 532nm wavelength.

Glutathione was estimated using 40µL of the reaction mix and summated with the specimen, positive control and reagent control wells, properly combined and was incubated at room temperature for 15 minutes to deplete all glutathione disulphite (GSSG) in the samples. 10µL of cumene hydroxide solution was added to start the glutathione peroxidase reaction and mixed well. The output (A1) was measured on a microplate reader at OD340nm at T1 and incubated at 25°C for 5 minutes (protected from light). The output (A2) was then measured on a microplate reader at OD340nm at T2.

Statistical analysis: Data were analyzed using a statistical package for social science version 21.0 The student t-test was used with data represented as mean ± standard error of the mean (SEM) and statistically significant at p < 0.05.

3. RESULTS

SHB administered to array II elevated MDA level (672 ± 39.65) when analogized to the negative

control array I (336 ± 39.21) and Donepezil treated array III (446 ± 78.33). TO (875mg/kg and 1750 mg/kg) administration to arrays IV and V meaningfully decreased MDA level (487 ± 78.33; 477 ± 29.50) when compared to SHB array II and TT (875mg/kg and 1750 mg/kg BW) treated arrays VI and VII (707.67 ±24.99; 671.64 ±32.07) at P<0.05 (Fig. 1). Rats in TT arrays VI and VII revealed minimal elevation compared to the SHB array II. Array V given an elevated dose of TO shows a meaningful reduction (P<0.05) in MDA levels as analogized to negative control array I (Fig. 1).

Increased glutathione is a label of cellular antioxidant and gives defense against oxidative

trouble. SHB treated array II rats (0.15±0.56), TT treated array VII (0.22±0.29) and Donepezil treated array III (0.20±0.00) revealed a meaningful declined glutathione status when analogized to control array I (0.49±0.21), TO treated arrays IV and V (0.41±0.08 and 0.46±0.07) and low dose TT treated array VI (0.44±0.05). The rats in array V treated with 1750mg/kg of TO revealed a slight reduction (P<0.05) of GSH levels analogized to the array I and significantly increased compared to Donepezil treated array III (0.46±0.18) and SHB treated array II. Rats in array VII treated with high dose TT (1750 mg/kg) showed a meaningful decrease (0.22±0.29) at P>0.05 (Fig. 2).

Table 1. Showing Malondialdehyde (MDA) and Glutathione (GPX) concentrations in diverse experimental arrays

Arrays	MDA (Mean ± SEM)	GSH (Mean ± SEM)
I	336±39.21	0.49±0.21
II	672±39.65	0.15±0.56
III	446±78.33	0.20±0.00
IV	487±78.33	0.41±0.08
V	477±29.50	0.46±0.07
VI	707.61±24.99	0.44±0.18
VII	671.64±32.07	0.22±0.29

Data are represented as mean±SEM, n=6.

* = Meaningfully dissimilar control at P < 0.05

a = Meaningfully dissimilar scopolamine hydrobromide at P < 0.05.




Fig. 1. Malondialdehyde concentrations in diverse experimental arrays

Data are represented as mean ± SEM, n=6.

* = Meaningfully dissimilar control at P < 0.05

a = Meaningfully dissimilar scopolamine hydrobromide at P < 0.05




Fig. 2. Glutathione (GPX) activity in the dissimilar experimental arrays

Data are represented as mean \pm SEM, n=6

* = Meaningfully dissimilar control at $P < 0.05$

a = Meaningfully dissimilar scopolamine hydrobromide at $P < 0.05$




Fig. 3. Biosynthesis of glutathione

4. DISCUSSION

The generation of free radicals associated with enlarged oxidative stress is attributed to the

pathogenesis of Alzheimer's disease resulting in ageing and cell apoptosis [16]. Studies suggested the involvement of oxidants in the neurological and neurodegradative ailments, thus

directing the way of free radicals in progression of Alzheimer's ailment. The use of herbal and natural extracts for neurological, psychiatric and neurotoxicological ailments has been elevated greatly because of their no or reduced side effects [17]. *Citrus esculentus* extract reversed radiation-induced damage on the ultrastructure of albino Wistar rats' testes [18], *Averrhoa carambola* aqueous fruit extract ameliorates diazepam-caused poison of the hippocampus in rats [19], *Musa paradisiaca* stem juice limit the extent of status epilepticus, increased neuronal protein synthesis, reduced cytoarchitectural damage and astriogliosis as well as enhancing long term recognition memory in rats [20,21] while *Ziziphus jujuba* fruit protects oxidative brain damage in rats [22]. Also, while Agbor and Anyanwu reported that β -D-Glucagon polysaccharide supplementation ameliorates hypoglycaemic-induced nephrotoxicity in diabetic Wistar rats [23], Cennet *et al* linked bulbs of *Muscari muscarimi* extract having more marked anticancer activities against H1299 cell line than other cell lines [24] and Nema *et al* demonstrated that erythropoietin is cytoprotective and blocked neurodegeneration by inhibiting caspase activity and apoptosis [25]; *Talinum triangulare* leaves possess antioxidant activities which was linked to amelioration of the atrophied hippocampal microstructures as well as aiding learning and recall in Alzheimer's type cognitive dysfunction rats [26].

In this study, there was a meaningful elevation of serum MDA levels in arrays II (treated with a SHB alone), VI and VII (treated with TT) analogous to the negative control in array I with a reduction in MDA level in arrays III, IV and V (Fig. 1). This result is in line with research that documented that TO seeds reduce lipid peroxidation at lower doses [27]. The SHB has been found to increase oxidative stress and improve the antioxidative defense system [28,29]. This is similar to reports that SHB significantly increased MDA levels in rats compared to the control [30]. Lipid peroxidation is a major pointer of neuron degradation of the brain. Unlike other body envelopes, neuronal envelopes encompass a very elevated percentage of long chain polyunsaturated fatty acids and oxidants are continually formed during metabolism [31]. Lipids and proteins, the important anatomical and physiological constituents of the cell membranes are the mark of oxidative alteration by free radicals in neurons degrading ailments [32]. A broad fact documented on lipid peroxidation and protein

oxidation resulting to loss of envelop structure, a good agent to increase aging and age-related neurodegenerated diseases. Oxidative trouble has been enlaced in the pathogenesis of AD in humans [33]. This still lends credence to the fact that neurodegenerative changes in histological, histochemical, immunohistochemical and neurobehavioral observations of the hippocampus could be attributed to activities of free radicals generated by SHB; hence, aqueous extract of TO seeds and TT leaves were able to ameliorate the effects [34-38].

Glutathione plays an important part in many metabolic and biochemical reactions such as DNA mixture and repair, protein mixture, prostaglandin mixture, amino acid movement and enzyme actuation [39]. According to Penninckx [40], glutathione also called γ -L-glutamyl-L-cysteinylglycerine is a redox-action tripeptide thiol found in the cells of a living organism. This bio-molecule serves as a detoxicant and acts as an antioxidant. During acute oxidative stress, the cells may not have the ability to reduce glutathione disulphite (GSSG) to GSH which may lead to accumulation of GSSG in the cell. This may be averted either by ingestion of diet containing antioxidants (such as TO seeds and TT leaves) or removing the accumulated GSSG from the cytosol. Hence, all system in the body can be influenced by the state of the glutathione system including the nervous system. In this study, a low level of glutathione was observed in arrays II, III and VII analogous to the negative control array I (Fig. 2). This correlates with the result from the MDA assay which showed increased levels of free radicals. This still showed a trend that scopolamine generates free radicals and increases oxidative stress (Fig. 1). A research work revealed that scopolamine caused behavioral and biochemical alterations in rats [41]. Their result showed that scopolamine-induced memory impairment increases MDA and decreases glutathione levels in hippocampal homogenate. Animals treated with TO seed and TT leaves showed increased levels of glutathione peroxidase compared to the SHB treated array II though the effect of TT was dose dependent (Fig. 2). The changes could be linked to the antioxidant capabilities of TO seeds and TT leaves. Consumption of antioxidants via diet and supplements is expected to turn off reactive oxygen species from the living system and give health benefits. The antioxidant potentials of TO seeds observed in biochemical analysis correlate with the Morris water maze test where extract of TO seeds aid learning and recall [36].

Glutathione is important antioxidant according to Lu [42] where free radicals (such as hydrogen peroxide) generated as a result of aerobic metabolism are metabolized by GSH peroxidase in the cytoplasm and mitochondria and by catalase in the peroxisome of a cell. The formed GSSG is reduced back to GSH by GSSG reductase at the expense of NADPH (Fig. 3). The organic peroxidase (ROOH) can be reduced by GSH peroxidase or GSH S-transferase. Under acute oxidative stress as seen in array II only (Fig. 2) may result to the inability of the cell to reduce GSSG to GSH, hence causing GSSG accumulation in the cell. To avoid a shift in the redox equilibrium, GSSG can be actively transported out of the cell or react with a protein sulfhydryl (PSH) to form a mixed disulfide (PSSG) (Fig. 3) [42]. In all the analysis, there is still a trend of protection observed that could be due to the plants' antioxidant potentials that may activate the biosynthesis of the glutathione leading to either the transport of GSSG out of the cell or forming PSSG by reacting with PSH.

5. CONCLUSION

In conclusion, aqueous extracts of TO seeds reduced MDA, TT leaves increased MDA and increase glutathione levels which may provide an enabling environment for cell survival and behavior using SHB-induced Alzheimer's type cognitive dysfunction rats.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Yin H, Xu L, Porter NA. Free radical lipid peroxidation: mechanisms and analysis. *Chem Rev.* 2011; 111 (10): 5944–5972.
2. Volinsky R, Kinnunen PKJ. Oxidized phosphatidylcholines in membrane-level cellular signalling: from biophysics to physiology and molecular pathology. *Fed Eur Biochem So J.* 2003; 280 (12) 2806–2816.
3. Mari M, Morales A, Colell A, Garcia-Ruiz C, Fernandez-Checa JC. Mitochondrial glutathione, a key survival antioxidant. *Ant Red Sig.* 2009; 11(11): 2685–2700.
4. Filomeni G, Aquilano K., Civitareale P, Rotillio G, Ciriolo MR. Activation of c-Jun-N-terminal kinase is required for apoptosis triggered by glutathione disulfide in neuroblastoma cells. *Free Radic Biol Med.* 2005; 39(3):345–354.
5. Ballatori N, Krance SM, Notenboom S, Shi S, Tieu K, Hammond CL. Glutathione dysregulation and the etiology and progression of human diseases. *J Biol Chem.* 2009; 390 (3): 191–214.
6. Su B, Wang X, Nunomura A, Moreira PI, Lee H, Perry G, et al. Oxidative stress signalling in Alzheimer's disease. *Cur Alzh Res.* 2008; 525-532.
7. Lian H, Yang L, Cole A, Sun L, Chiang ACA, Fowler SW, et al. NFKB-activated astroglial release of complement C3 comprises neuronal morphology and function associated with Alzheimer's disease. *Neu.* 2015; 85, 101-115.
8. Oda A, Tamaoka A, Araki W. Oxidative stress up-regulates presenilin 1 in lipid rafts in neuronal cells. *J Neurosci Res.* 2010; 88(5): 1137–1145.
9. Yankner BA, Lu T. Amyloid β -protein toxicity and the pathogenesis of Alzheimer disease. *J Bio Chem.* 2009; 284(8): 4755–4759.
10. Manczak M, Anekonda TS, Henson E, Park BS, Quinn J, Reddy PH. Mitochondria are a direct site of A β accumulation in Alzheimer's disease neurons: implications for free radical generation and oxidative damage in disease progression. *Hum Mol Gen.* 2006;15(9):1437-1449.
11. Muhammad-Abdul H, Sultana R, Keller JN, St. Clair DK, Markesbery WR, Butterfield DA. Mutations in amyloid precursor protein and presenilin-1 genes increase the basal oxidative stress in murine neuronal cells and lead to increased sensitivity to oxidative stress mediated by amyloid β -peptide (1-42), H₂O₂ and kainic acid: implications for Alzheimer's disease. *J Neurochem.* 2006;96(5):1322-1335.
12. De Felice FG, Velasco PT, Lambert MP. A β oligomers induce neuronal oxidative stress through an N-methyl-D-aspartate

- receptor-dependent mechanism that is blocked by the Alzheimer drug memantine. *J Bio Chem.* 2007; 282(15): 11590–11601.
13. Zhang L, Ravipati AS, Koyyalamudi SR, Jeong SC, Reddy N, Smith PT, et al. Antioxidant and antiinflammatory activities of selected medicinal plants containing phenolic and flavonoid compounds. *J Agric Food Chem.* 2011;59(23):12361-7.
 14. Pharm-Huy LA, He H, Phar-Huy C. Free radical. Antioxidants in disease and health. *Int J Biomed Sci.* 2008;4 (2):89-96.
 15. Lorke D. A new approach to practical acute toxicity testing. *Archives of Toxicology.* 1983;54: 275-287.
 16. Valko M, Leifritz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *The Int J Bioch Cell Bio.* 2007;39:44-84.
 17. Kim SJ, Lee JH, Chung HS, Song JH, Ha J, Bae H. Neuroprotective Effects of AMP-activated protein kinase on scopolamine induced memory impairment. *Korean J Phy Pharm.* 2013;17: 331-338.
 18. Udoh BE, Erim AE, Paulinus SO, Eru EM, Ukpong EV, Efang I, et al. Effects of *Cyperus esculentus* (Tiger Nut) Extract on the Irradiated Testes of Wistar Rats. *Trop J Nat Prod Res.* 2020; 4(11):966-969.
 19. Anani SE, Eru EM, Okon DE, Uruakpa KC, Ugbem TI, Igiri AO. The Effect of *Averrhoa carambola* (Star Fruit) Aqueous Fruit Extract on the Hippocampal Astrocyte Expression following Diazepam-Induced Neurotoxicity in Wistar Rats. *Trop J Nat Prod Res.* 2020;4(12):1170-1173.
 20. Ifio FB, Eru EM, Mathias OA, Nsikak MU, Williams AN, Florence NO et al. Effect of musa paradisiaca stem juice on acute status epilepticus, hippocampal histology and behaviour in pentylenetetrazole – induced wistar rats. *Niger. J. Physiol. Sci.* 2024;39: (in press).
 21. Ifio FB, Gabriel UU, Eru EM, Nsikak MU, Mathias OA, Williams AN et al. Histochemical study of Nissl substance and astrocytes in a pentylenetetrazole-induced model of epilepsy treated with musa paradisiaca stem juice. *Niger. J. Physiol. Sci.* 2024;39: (in press).
 22. Mohammed SA. Effects of ziziphus jujuba fruits extract on memory impairment induced hypothyroidism during breastfeeding and adolescence in the rats. *Jordan J Biol Sci.* 2021;15 (1):119-125.
 23. Agbor CA, Anyanwu GE. Antioxidant effect of beta-D-glucagon-polysaccharide fractionate of *Auricularia polytricha* on hyperglycaemia-induced kidney dysfunction in experimental diabetic nephropathy. *Jordan J Biol Sci.* 2021; 5(14):905-910.
 24. Cennet O, Ege RK, Ramazan M, Hakan A. Antioxidant and apoptotic effect of muscari muscarimi, an endemic geophytes species from turkey. *Jordan J Biol Sci.* 2021; 4(14):819-823.
 25. Nema AM, Mona IS, Amal LE, Hussein KHH, Trez NM. The ameliorating effect of erythropoietin on diabetic neurodegeneration by modulating the antioxidant-oxidant imbalance and apoptosis in diabetic male rats. *Jordan J Bio Sci.* 2018; 3(11): 339-345.
 26. Eru EM, Paulinus SO, Igiri AO, Akpaso MI. Enhanced effect of aqueous extract of *Telfairia occidentalis* seed on the microstructure of the hippocampus of scopolamine hydrobromide-induced cognitive dysfunction rats. *Asian J of Res in Neuro.* 2020;3(1):5-10.
 27. Oro-Oluwapo OD, Wahab AO, Gideon O. Effects of methanol extract of *Telfairia occidentalis* seed on serum lipid profile, biochemical and antioxidant activity in female wistar rats. *Eur J Med Pl.* 2016;15(2): 1-8.
 28. Ishola IO, Tota S, Adeyemi OO, Agbaje EO, Narender T, Shukla R. Protective effect of *Cnestis feruginea* and its active constituent on scopolamine-induced memory impairment in mice: a behsvioral and biochemical study. *Pharm Bio.* 2013; 51:825-835.
 29. Marisco PC, Carvalho FB, Rosa MM, Girardi BA, Gutierrez JM, Jaques JA. Piracetam prevents scopolamine-induced memory impairment and decrease of NTPDase, 5'-nucleotide and adenosine deaminase activities. *Neurochem Res.* 2003; 38:1704-1714.
 30. Kaur R, Mehan S, Khanna D, Kalra S. Ameliorative Treatment with Ellagic Acid in Scopolamine Induced Alzheimer's Type Memory and Cognitive Dysfunctions in Rats. *Austin J Clin Neu.* 2015;2 (6):1053-1064.
 31. Gella A, Durany N. Oxidative stress in Alzheimer disease. *Cell Ad Migr.* 2009;3:88-93.
 32. Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional

- foods: Impact on human health. Pharm Rev. 2010;4:118- 126.
33. Uttara B, Singh AV, Zamboni P, Mahajan RT. Oxidative stress and neurodegenerative diseases: A review of upstream and downstream antioxidant therapeutic options. Cur Neuropharm. 2009;7:65-74.
 34. Eru EM, Anani SE, Paulinus SO, Mesembe OE, Uruakpa KC, Umoh NM, et al. Scopolamine-Induced Alzheimer's Disease in Wistar Rats; Aqueous *Talinum triangulare* Potency on the Hippocampal Nissl Bodies and Long-Term Learning and Memory. Trop J Nat Prod Res. 2022; 6(1):117-122.
 35. Eru ME, Gabriel UU, Samson OP, Kelechi CU, Michael EO, Sadeyeng EA, et al. Restorative Potentials of Aqueous *Telfairia occidentalis* Seeds Extract on the Hippocampal Nissl Granules and Short-Term Memory in Scopolamine Hydrobromide-Induced Alzheimer's Type Cognitive Dysfunction Rats. Trop J Nat Prod Res. 2021; 5(1):182-187
 36. Eru EM, Paulinus SO, Udo-Affah GU, Uruakpa KC, Oku ME, Anani SE et al. Hippocampal Astroglial Reduction in Scopolamine Hydrobromide-Induced Alzheimer's Cognitive Dysfunction Wistar Rats Following Administration of Aqueous Extract of *Telfairia occidentalis* (Hook F.) Seeds. Niger J Physiol Sci, 2021; 36: 241 – 244
 37. Eru EM, Paulinus SO, Gabriel UU, Kelechi CU, Michael EO, Edet UI, et al. Neurobehavioural enhancement in scopolamine hydrobromide-induced alzheimer type cognitive dysfunction in rats following administration of ethanol seed extract of *Telfairia occidentalis* (hook.f.) cucurbitaceae. Trop J Nat Prod Res, 2020; 4(7):282-285.
 38. Eru EM, Gabriel UU, Ifioke FB, Kelechi CU, Samson OP, Michael EO et al. Efficacy of aqueous extract of *Talinum triangulare* on the microanatomy of the hippocampus and short-term memory of scopolamine hydrobromide-induced alzheimer's type cognitive dysfunction rats. Niger. J. Physiol. Sci. 2024;39 (in press).
 39. Allen J, Bradley RD. Effects of oral glutathione supplementation on systemic oxidative stress biomarkers in human volunteers. J Alt Comp Med. 2011;17(9): 827-833.
 40. Penninckx MJ, Elsken MT. Metabolism and functions of glutathione in micro-organism. Adv Micro Physiol. 1993;34:239-301.
 41. Lu SC. Regulation of hepatic glutathione synthesis: current concept and controversies. FASEB J. 1999;13 (10):1169-1183.
 42. Puri A, Srivastava P, Pandey P, Yadav RJ, Bhatt PC. Scopolamine induced behavioural and biochemical modifications and protective effect of *Celastrus paniculatus* and *Angelica glauca* in rats. Int J Nut Pharm Neu Dis. 2014; 4(3): 158-169.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/121691>