



Seasonal Variation in an Outpatient Antibiotic Prescription Rates in Alkharj

Nehad J. Ahmed^{1*}

¹*Department of Clinical Pharmacy, College of Pharmacy, Prince Sattam Bin Abdulaziz University, Al-kharj, Saudi Arabia.*

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/JPRI/2020/v32i930476

Editor(s):

(1) Vasudevan Mani, Qassim University, Saudi Arabia.

Reviewers:

(1) Fils Landry Mpelle, Marien Ngouabi University, Republic of the Congo.

(2) Ruqiya Sultana, Omega BIBI Hospital, India.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/55728>

Original Research Article

Received 24 January 2020

Accepted 30 March 2020

Published 03 June 2020

ABSTRACT

Aim: The aim of this study was to describe the seasonal variations in dispensing antibiotic prescriptions in the outpatient setting of a public hospital in Alkharj.

Methodology: A retrospective cross-sectional study was conducted. The outpatient prescriptions in 2017 and 2018 were collected from medical records in a public hospital in Alkharj.

The data include the number of prescribed antibiotics in general, the number of prescribed antibiotics in different months and seasons in the outpatient setting.

Results: In the outpatient setting in 2017 and 2018, antibiotics were prescribed excessively in most of the months. About 27.84% of the prescriptions in 2017 were in spring season and about 26.64% of the prescriptions in 2018 were in autumn. Total number of antibiotics prescriptions in 2017 and 2018 were 5348 in spring followed by 5097 in autumn.

Conclusion: The results of the present study showed the widespread use of antibiotics by practitioners that was associated with season of prescribing. In general, there are excess use of antibiotics in all months. It is important to understand how the prescribing of antibiotic varies throughout the year to design an appropriate intervention to decrease incorrect antibiotic use.

Keywords: Seasonal variation; outpatient; out-patients; antibiotics antibacterial agents.

*Corresponding author: E-mail: n.ahmed@psau.edu.sa, pharmdnehadjaser@yahoo.com;

1. INTRODUCTION

Antibiotics are drugs that are used to treat and to prevent bacterial infections but not for other infections such as viral infections. Antibiotic resistance occurs when bacteria change in response to the usage of these medicines. Bacteria become frequently resistant to antibiotic. These bacteria may infect not only humans but also animals, and the infections they cause are harder to treat than those caused by non-resistant bacteria [1].

Antibiotic resistance continues to be a threat that affects the efficacy of antibiotics and poses a significant challenge for prescribers who treat patients with bacterial infections. Infections caused by resistant organisms limit the clinicians' management options because there is a lack in the development and availability of novel antibiotics [2,3]. There is a growing evidence from previous studies that the excessive use of antibiotics is a main risk factor for the development of antibiotic resistance [4,5]. Therefore, to decrease the number of infections caused by resistant bacteria, unsuitable antibiotic prescribing must be decreased [6-11].

In the United States the majority of antibiotics are prescribed in primary care settings for the management of infections in the upper respiratory tract, for which antibiotic therapy is occasionally needed [12-17].

Additionally, most of the antibiotics are prescribed in the first and fourth quarters of the calendar year particularly in winter, because in winter the bacterial and viral illnesses such as influenza and rhinosinusitis are common [18-20].

It is important to identify when and where to use appropriate interventions to decrease antibiotic use in ambulatory care. The aim of this study was to describe the seasonal variations in dispensing antibiotic prescriptions in the outpatient setting of a public hospital in Alkharj in order to identify the opportunity for clinical interventions where seasonal peaks may show increases in unsuitable antibiotic use.

2. METHODOLOGY

A retrospective cross-sectional study was conducted. The outpatient prescriptions in 2017 and 2018 were collected from medical records in a public hospital in Alkharj.

The data include the number of prescribed antibiotics in general, the number of prescribed antibiotics in different months and seasons in the outpatient setting. We included only the most commonly used antibiotics which are amoxicillin / clavulanic acid, amoxicillin, metronidazole, ciprofloxacin, cefuroxime and azithromycin.

The exclusion criteria include the inpatient prescriptions and the outpatient prescription before 2017 and after 2018, and other antibiotics that were not prescribed commonly.

The data were collected and analyzed using excel software. The data represented by frequencies and percentages. This study is approved by Institutional Review Board Log No. 18-474E.

3. RESULTS

In the outpatient setting in 2017 and 2018, antibiotics were prescribed excessively in most of the months. Table 1 shows the outpatient prescribing of the most common antibiotics in different months in 2017 and Table 2 shows the outpatient prescribing of the most common antibiotics in different months in 2018.

About 27.84% of the prescriptions in 2017 were in spring season and about 26.64% of the prescriptions in 2018 were in autumn. Table 3 shows the outpatient prescribing of the most common antibiotics in different seasons in 2017 and Table 4 shows the outpatient prescribing of the most common antibiotics in different seasons in 2018.

Total number of antibiotics prescriptions in 2017 and 2018 were 5348 in spring followed by 5097 in autumn, 4966 in winter and 4409 in summer. The outpatient prescribing of the most common antibiotics in different seasons in 2017 and 2018 is shown in Table 5 and the total prescriptions in different months are shown in Table 6.

4. DISCUSSION

A high percentage of the prescriptions in 2017 were in spring season (27.84%) and in 2018 a high percentage of the prescriptions in 2018 were in autumn (about 26.64%). For the total antibiotics use for 2 years in the present study, spring followed by autumn were the seasons of the highest prescriptions more than winter and summer.

Table 1. The outpatient prescribing of the most common antibiotics in different months in 2017

Antibiotic	March	April	May	June	July	Aug	Sep	Oct	Nov	Dec	Jan	Feb
Metronidazole	77	76	70	82	105	81	57	75	62	57	79	75
Ciprofloxacin	68	78	66	41	56	46	63	80	51	51	53	40
Cefuroxime	74	87	57	38	101	58	70	54	40	38	21	49
Amoxicillin	367	304	282	208	334	325	328	389	272	301	364	281
Amoxicillin/ Clavulanic acid	211	318	269	225	227	189	154	242	218	227	246	150
Azithromycin	91	85	138	105	76	31	106	86	124	132	81	54
Total	888	948	882	699	899	730	778	926	767	806	844	649

Table 2. The outpatient prescribing of the most common antibiotics in different months in 2018

Antibiotic	March	April	May	June	July	Aug	Sep	Oct	Nov	Dec	Jan	Feb
Metronidazole	133	139	97	81	88	103	96	81	86	86	76	96
Ciprofloxacin	59	60	55	62	66	62	62	78	67	69	60	73
Cefuroxime	53	56	38	35	43	27	25	43	42	76	31	24
Amoxicillin	306	265	202	270	231	225	321	279	299	395	312	290
Amoxicillin / Clavulanic acid	315	306	273	169	156	208	233	262	336	223	350	257
Azithromycin	82	116	75	86	78	91	118	116	136	67	79	103
Total	948	942	740	703	662	716	855	859	966	916	908	843

Table 3. The outpatient prescribing of the most common antibiotics in different seasons in 2017

Season	Number of antibiotics prescribed	Percentage of antibiotics prescribed
Spring	2718	27.84
Summer	2328	23.84
Autumn	2417	24.75
Winter	2299	23.55

Table 4. The outpatient prescribing of the most common antibiotics in different seasons in 2018

Season	Number of antibiotics prescribed	Percentage of antibiotics prescribed
Spring	2630	26.14
Summer	2081	20.68
Autumn	2680	26.64
Winter	2667	26.51

Table 5. The outpatient prescribing of the most common antibiotics in different seasons in 2017 and 2018

Variable	Spring	Summer	Autumn	Winter
Number of antibiotics prescribed in 2017 and 2018 (Total)	5348	4409	5097	4966
Percentage of total antibiotics prescribed	26.98	22.24	25.71	25.05

Katie J. Suda et al. [21] reported that the antibiotic prescribing is higher in winter than in summer but in their study they compare between the prescriptions in 2 seasons only summer and winter. Similar to this result in our study the prescriptions in winter were higher than in summer but less than autumn and spring.

In contrast to our study, the antibiotic prescriptions in winter are more than other seasons as reported by several European and Canadian studies, they stated that the increase in antibiotics use in the winter months ranged from 21% to 42% [19,22-24]. Moreover, Bauchner H et al. reported that antibiotics are

Table 6. The total outpatient prescribing of the most common antibiotics in different months

Antibiotic	Marh	April	May	June	Juy	Aug	Sep	Oct	Nov	Dec	Jan	Feb
Prescriptions in 2017	888	948	882	699	899	730	778	926	767	806	844	649
Prescriptions in 2018	948	942	740	703	662	716	855	859	966	916	908	843
Total Prescriptions	1836	1890	1622	1402	1561	1446	1633	1785	1733	1722	1752	1492

prescribed in winter more than other seasons and that a high percentage of antibiotic prescriptions in the winter are estimated to be unsuitable [25].

Additionally, previous study stated that the use of antibiotic is more in winter and that the excess use of antibiotics in the winter could have a significant effect on resistance [18].

Many physicians prescribe antibiotics to treat viral infections. Seasonal peaks in antibiotics generally used to treat viral upper respiratory tract infections remained unchanged during cold and influenza season [26].

Seasonal influenza viruses are detected year-round, but viruses are most common during the fall and winter. The timing and duration of flu seasons can vary, but generally influenza activity begins to increase in October. Most of the time flu activity peaks between December and February, although activity can last as late as May [27]. In the present study, the highest prescribing was in April followed by March.

Pathak A et al reported that in the peak-prescribing rate of antibiotics was during the summer (75%), while the teaching hospital had a peak prescribing rate during the rainy season (70%) and this study also in contrast to our study [28].

It is noticed that the seasonal variations in prescribing antibiotics are different for the various antibiotics. Highest prescribing rate of metronidazole (a nitroimidazole antibiotic) was in April followed by march, highest prescribing rate of ciprofloxacin (a fluoroquinolone antibiotic) was in October followed by April, highest prescribing rate of cefuroxime (a cephalosporin antibiotic) was in July followed by April, highest prescribing rate of amoxicillin (a penicillin antibiotic) was mainly in December followed by March, highest prescribing rate of amoxicillin/clavulanic acid (a penicillin antibiotic) was mainly in April followed by January and highest prescribing rate of azithromycin (macrolide antibiotic) was in November followed by September.

Amoxicillin/clavulanic acid, metronidazole and cefuroxime were prescribed mostly in spring, ciprofloxacin and azithromycin were prescribed mainly in autumn, amoxicillin was prescribed mainly in winter.

In contrast to our study, Safaeian L, et al. stated that that seasonal peak was observed for

penicillins and cephalosporins prescriptions in autumn. But regarding macrolides, similarly he reported that they were significantly more prescribed during autumn [29]. Furthermore, Suda KJ et al. reported that more antibiotic prescribing, predominately driven by the macrolide and penicillin classes, in the outpatient setting was observed in the winter months [21].

5. CONCLUSION

The results of the present study showed the widespread use of antibiotics by practitioners that was associated with season of prescribing. Amoxicillin and amoxicillin/clavulanic acid were the most prescribed antibiotics. In general, there were excess use of antibiotics in all months, but each antibiotic was prescribed more in specific months. It is important to understand how the prescribing of antibiotic varies throughout the year; this will be useful for designing an appropriate intervention strategy to decrease incorrect antibiotic use. Antibiotics should be use as per antibiotic stewardship in order to promote rational use of antibiotics.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

ACKNOWLEDGEMENT

This Publication was supported by the Deanship of Scientific Research at Prince Sattam bin Abdulaziz University.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. World Health Organization. Fact sheet–antibiotic resistance–media center. World Health Organization (WHO); 2016. Available: <https://www.WHO.int/en/news-room/fact-sheets/detail/antibiotic-resistance> (Accessed October, 10, 2019)
2. Spellberg B, Guidos R, Gilbert D, Bradley J, Boucher HW, Scheld WM, Bartlett JG,

- Edward J, Jr, Infectious Diseases Society of America. The epidemic of antibiotic-resistant infections: A call to action for the medical community from the Infectious Diseases Society of America. *Clin. Infect. Dis.* 2008;46:155–164.
3. Boucher HW, Talbot GH, Benjamin DK, Bradley J, Guidos RJ, Jones RN, Murray BE, Bonomo RA, Gilbert D, Infectious Diseases Society of America. 10 x '20 progress — development of new drugs active against gram-negative bacilli: An update from the Infectious Diseases Society of America. *Clin. Infect. Dis.* 2013; 56:1685–1694.
 4. Seppälä H, Klaukka T, Vuopio-Varkila J, Muotiala A, Helenius H, Lager K, et al. The effect of changes in the consumption of macrolide antibiotics on erythromycin resistance in group A streptococci in Finland. *New Engl J Med.* 1997;337:441–446.
 5. Hay AD, Thomas M, Montgomery A, Wetherell M, Lovering A, McNulty Cet al. The relationship between primary care antibiotic prescribing and bacterial resistance in adults in the community: a controlled observational study using individual patient data. *J Antimicrob Chemother.* 2005;56:146–153.
 6. Austin DJ, Kristinsson KG, Anderson RM. The relationship between the volume of antimicrobial consumption in human communities and the frequency of resistance. *Proc. Natl. Acad. Sci. U. S. A.* 1999;96:1152–1156.
 7. Goossens H, Ferech M, Vander Stichele R, Elseviers M; ESAC Project group. Outpatient antibiotic use in Europe and association with resistance: Across-national data base study. *Lancet.* 2005; 365:579–587.
 8. Hicks LA, Chien YW, Taylor TH, Haber M, Klugman KP, Active Bacterial Core Surveillance (ABCs) Team. Outpatient antibiotic prescribing and non-susceptible *Streptococcus pneumoniae* in the United States, 1996–2003. *Clin. Infect. Dis.* 2011; 53:631–639.
 9. Jensen US, Muller A, Brandt CT, Frimodt-Moller N, Hammerum AM, Monnet DL, DANRES study group. Effect of generics on price and consumption of ciprofloxacin in primary healthcare: The relationship to increasing resistance. *J. Antimicrob. Chemother.* 2010;65:1286–1291.
 10. Malhotra-Kumar S, Lammens C, Coenen S, Van Herck K, Goossens H. Effect of azithromycin and clarithromycin therapy on pharyngeal carriage of macrolide-resistant streptococci in healthy volunteers: a randomized, double-blind, placebo-controlled study. *Lancet.* 2007;369:482–490.
 11. Metz-Gercek S, Maieron A, Strauss R, Wieninger P, Apfalter P, Mittermayer H. Ten years of antibiotic consumption in ambulatory care: Trends in prescribing practice and antibiotic resistance in Austria. *BMC Infect. Dis.* 2009;9:61.
 12. McCaig LF, Hicks LA, Roberts RM, Fairlie TA. Office-related antibiotic prescribing for persons aged 14 years—United States, 1993–1994 to 2007–2008. *MMWR Morb. Mortal. Wkly. Rep.* 2011;60:1153–1156.
 13. Hersh AL, Shapiro DJ, Pavia AT, Shah SS. 2011. Antibiotic prescribing in ambulatory pediatrics in the United States. *Pediatrics.* 2011;128:1053–1061.
 14. Grijalva CG, Nuorti JP, Griffin MR. Antibiotic prescription rates for acute respiratory tract infections in US ambulatory settings. *JAMA.* 2009;302:758–766.
 15. McCaig LF, Besser RE, Hughes JM. Antimicrobial drug prescription in ambulatory care settings, United States, 1992–2000. *Emerg. Infect. Dis.* 2003;9: 432–437.
 16. Steinman MA, Gonzales R, Linder JA, Landefeld CS. Changing use of antibiotics in community-based outpatient practice, 1991–1999. *Ann. Intern. Med.* 2003;138: 525–533.
 17. Fairlie T, Shapiro DJ, Hersh AL, Hicks LA. National trends in visit rates and antibiotic prescribing for adults with acute sinusitis. *Arch. Intern. Med.* 2012;172:1513–1514.
 18. Sun L, Klein EY, Laxminarayan R. Seasonality and temporal correlation between community antibiotic use and resistance in the United States. *Clin. Infect. Dis.* 2012;55:687–694.
 19. Elseviers MM, Ferech M, Vander Stichele RH, Goossens H, ESAC Project Group. Antibiotic use in ambulatory care in Europe (ESAC data 1997–2002): trends, regional differences and seasonal fluctuations. *Pharmacoepidemiol. Drug Saf.* 2007;16: 115–123.
 20. Ferech M, Coenen S, Malhotra-Kumar S, Dvorakova K, Hendrickx E, Suetens Cet al. European surveillance of antimicrobial

- consumption (ESAC): outpatient antibiotic use in Europe. *J. Antimicrob. Chemother.* 2006;58:401–407.
21. Suda KJ, Hicks LA, Roberts RM, Hunkler RJ, Taylor TH. Trends and seasonal variation in outpatient antibiotic prescription rates in the United States, 2006 to 2010. *Antimicrob Agents Chemother.* 2014;58(5):2763-6.
 22. Achermann R, Suter K, Kronenberg A, Gyger P, Muhlemann K, Zimmerli W, et al. Antibiotic use in adult outpatients in Switzerland in relation to regions, seasonality and point of care tests. *Clin. Microbiol. Infect.* 2011;17:855–861.
 23. Patrick DM, Marra F, Hutchinson J, Monnet DL, Ng H, Bowie WR. Per capita antibiotic consumption: How does a North American jurisdiction compare with Europe? *Clin. Infect. Dis.* 2004;39:11–17.
 24. Dagan R, Bakai G, Givon-Lavi N, Sharf AZ, Vardy D, Cohen T, et al. Seasonality of antibiotic-resistant *Streptococcus pneumoniae* that causes acute otitis media: A clue for an antibiotic restriction policy? *J. Infect. Dis.* 2008;197:1094–1102.
 25. Bauchner H, Pelton SI, Klein JO. Parents, physicians and antibiotic use. *Pediatrics.* 1999;103:395–401.
 26. Durkin MJ, Jafarzadeh SR, Hsueh K, Sallah YH, Munshi KD, Henderson RR, et al. Outpatient antibiotic prescription trends in the United States: A National Cohort Study. *Infect Control Hosp Epidemiol.* 2018;39(5):584-589.
 27. Centers for Disease Control and Prevention. The Flu Season; 2018. Available: <https://www.cdc.gov/flu/about/season/flu-season.htm> [Accessed 32020].
 28. Pathak A, Mahadik K, Dhaneria SP, Sharma A, Eriksson B, Lundborg CS. Antibiotic prescribing in outpatients: Hospital and seasonal variations in Ujjain, India. *Scand J Infect Dis.* 2011;43(6-7): 479-88.
 29. Safaeian L, Mahdanian AR, Salami S, Pakmehr F, Mansourian M. Seasonality and physician-related factors associated with antibiotic prescribing: A cross-sectional study in Isfahan, Iran. *Int J Prev Med.* 2015;15;6:1.

© 2020 Ahmed; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/55728>