



# Effect of Nitrogen and Spacing on Green Fodder Yield and Economics of MP Chari (*Sorghum bicolor* L.)

Mudakala Lokesh Chand<sup>a+++\*</sup>, Rajesh Singh<sup>a#</sup>,  
Rathod Sandeep<sup>a++</sup>, Akhankya Pradhan<sup>a†</sup>,  
and Thakur Indu<sup>a†</sup>

<sup>a</sup> Department of Agronomy, Faculty of Agriculture, Naini Agriculture Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj - 211007, Uttar Pradesh, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i82881

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/98149>

Original Research Article

Received: 25/01/2023  
Accepted: 29/03/2023  
Published: 01/04/2023

## ABSTRACT

At the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj (U.P.), India, a field experiment was carried out during the Harvest season of 2022. To investigate how spacing and nitrogen affect sorghum growth and fodder production (*Sorghum bicolor* L.). The treatments include nitrogen at rates of 60, 80, and 100 kg/ha, spaced 35, 45, and 55 cm, respectively. There were 10 treatments total, with three replications of each. The soil in the trial plot had a sandy loamy texture, a pH of 7.8 that was almost neutral, a low level of organic carbon (0.35%), and high levels of available N (163.42 kg/ha), available P (21.96 kg/ha), and available K (256.48 kg/ha) nutrients. In

<sup>++</sup> M. Sc Scholar;

<sup>#</sup> Associate Professor;

<sup>†</sup> PhD Scholar;

\*Corresponding author: E-mail: lokeshmudakala@gmail.com;

accordance with the results, nitrogen application at a rate of 100 kg per hectare and a spacing of 55 cm apart had a significant impact on plant height (186.96 cm), plant dry weight (52.78 g/plant), crop growth rate (55.0 g/m<sup>2</sup>/day), and green fodder yield (28.36 and 22.43 t/ha) for the first and second cuttings. With nitrogen 100 kg/ha and the spacing 5515cm, there were also larger gross returns (INR 83809.00/ha), net returns (INR 58133.00/ha), and B:C ratios (2.26).

**Keywords:** Sorghum; nitrogen; spacing; green fodder yield and economics.

## 1. INTRODUCTION

Sorghum (*Sorghum bicolor* (L.) Moench), the primary kharif and rabi season crop of the Poaceae family, is widely cultivated to meet the demands of livestock for both green and dry feed. The sudanese and bicolor are two species that are important economically. Bicolor is utilised as food, feed, and forage in India, whilst Sudanese are only used as fodder. All three crops—sorghum, Sudan grass, and hybrids of the two—are adaptable and have a range of uses. Grain sorghum is utilised for different purposes than forage sorghum. In many parts of the world, sorghum is evolving into a more major feed crop. It is superior to corn in terms of drought resistance and produces more with less irrigation, making it the ideal feed crop for desert regions. Sorghum can be used as fodder in a variety of ways, such as single-cut varieties for rain-fed farming, multi-cut varieties for irrigated agriculture in the form of stover, silage for storage, and other long-term uses.

For the purpose of generating assets and increasing livestock output, protein availability is a significant factor that affects animal productivity in terms of yield and quality. High protein intake may facilitate rapid weight gain and milk production [1].

The lack of acceptable genotypes and appropriate and correct fertilisation is one of the main factors limiting the production of fodder sorghum in our country [2]. For the production of fodder, nitrogen application is essential. According to Iqbal et al. [3], nitrogen supplementation increases the yield, dry matter production, and quality of oat-based green fodder. Furthermore, it was found by Ayub M. [4] and Saleem et al. [5] that the time and manner of nitrogen treatment significantly affected the yield of a number of maize hybrids. adequate mowing and the appropriate the two most important agronomic factors for generating higher fodder yields are planting geometry.

One of the essential elements of the crop establishment technique that influences the crop stand and other yield metrics in many crops is row spacing. A persistent problem for farmers is choosing the right planting density. Yields, radiation effectiveness, and weed infestation all suffer from lower plant density. However, a dense plant population may result in lodging, restricted light penetration in the canopy, a loss in photosynthesis due to lower leaves being shaded, and a markedly slowed pace of production [6].

Keeping these points in view, the present investigation entitled “Influence of Nitrogen and Spacing on Green fodder Yield and Economics of M P Chari (*Sorghum bicolor* L.)” was conducted during Zaid-2022, at crop research farm, SHUATS, Prayagraj (U.P.).

## 2. MATERIALS AND METHODS

During the zaid season of 2022, a field experiment was carried out at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.), India. The experimental plot's soil was sandy loamy in texture, nearly neutral in soil reaction (pH 7.8), and low in organic carbon (0.35%). The treatments included nitrogen at 60 kg per ha with spacing of 35 x 15 cm, 45 x 15 cm, and 55 x 15 cm, as well as nitrogen at 80 kg per ha with spacing of 35 x 15 cm, 100 kg per ha with spacing of 45 x 15 cm, and 100 kg per ha with spacing of 55 x 15 cm. Ten treatments were reproduced three times in a randomised block design experiment. Plant height, dry weight, the number of leaves per plant, crop growth rate (g/m<sup>2</sup>/day), and the yield of green fodder (t/ha) were all noted. The data was gathered and statistical analysis using the analysis of variance approach was performed [7].

### 3. RESULTS AND DISCUSSION

#### 3.1 Yield Attribute

**Green fodder Yield (t/ha):** The significantly higher green fodder yield (28.36 t/ha) was observed in treatment-9 (Nitrogen 100 kg/ha + Spacing 55x15cm) during 1<sup>st</sup> cutting. However, treatment-8 (Nitrogen 100 kg/ha + Spacing 45x15cm) was found to be statistically at par with treatment- 9 (Nitrogen 100 kg/ha + Spacing 55x15cm). The significantly higher green fodder yield (22.43 t/ha) was observed in treatment-9 (Nitrogen 100 kg/ha + Spacing 55x15cm) during 2<sup>nd</sup> cutting. However, treatment-8 (Nitrogen 100 kg/ha + Spacing 45x15cm) was found to be statistically at par with treatment- 9 (Nitrogen 100 kg/ha + Spacing 55x15cm). The significantly higher green fodder yield (28.36 t/ha) was observed with the application of nitrogen 100 kg/ha along with the spacing 55x15 cm, The increased plant height, number of leaves, plant population per metre, and leaf to stem ratio all contributed to the higher green fodder output at increasing spacing. In addition, the excessive plant population may fight with each other for light and nutrients, resulting in lanky growth and grassy shoot appearance, which reduced the output of green fodder at decreasing spacing. The benefits of nitrogen on cell division and

elongation, the formation of nucleotides and co-enzymes that resulted in an increase in meristematic activity and photosynthetic area and, consequently, more production and accumulation of photosynthates, yielding higher green fodder, are among the nitrogen levels that are primarily attributed to improved growth and yield parameters. These parameters include plant height, number of leaves, plant population per metre, leaf: stem ratio, plant height and dry weight. Similar results are conformity with Mane Shivprasad and Rajesh Singh [8], Dudhat et al. [9]. "The Optimum planting pattern is that the necessity for proper utilization of growth resources and ultimately to use the potential productivity of any crop. The higher grain yield was recorded from the interaction impact of sowing dates and spacing" Manasa and Umeha c. (2022). "The green fodder yield (28.36 t/ha) was observed with the application of nitrogen 100 kg/ha along with the spacing 55x15 cm, This may be with nitrogen mainly attributed to improved growth and yield parameters, viz., plant height, leaf area, leaf stem ratio and the beneficial effects of nitrogen on cell division and elongation, formation of nucleotides and Co-enzymes which resulted in increased meristematic activity and photosynthetic area and hence more production and accumulation of photosynthates, yielding higher green fodder" [4].

**Table 1. Influence of nitrogen and spacing on green forage yield of fodder sorghum M P Chari**

Treatment No.	Treatment combinations	Green fodder (t/ha) 1 <sup>st</sup> cutting	Green fodder (t/ha) 2 <sup>nd</sup> cutting
1.	Nitrogen 60 kg/ha + Spacing 35x15cm	22.68	18.17
2.	Nitrogen 60 kg/ha + Spacing 45x15cm	23.97	18.74
3.	Nitrogen 60 kg/ha + Spacing 55x15cm	25.09	19.31
4.	Nitrogen 80 kg/ha + Spacing 35x15cm	24.08	18.39
5.	Nitrogen 80 kg/ha + Spacing 45x15cm	25.19	19.52
6.	Nitrogen 80 kg/ha + Spacing 55x15cm	27.03	21.60
7.	Nitrogen 100 kg/ha + Spacing 35x15cm	26.41	20.08
8.	Nitrogen 100 kg/ha + Spacing 45x15cm	27.45	21.73
9.	Nitrogen 100 kg/ha + Spacing 55x15cm	28.36	22.43
10.	Control	23.25	18.85
F test		S	S
SEm±		0.71	0.34
CD (P=0.05)		2.10	1.02

**Table 2. Economic analysis of different treatment combinations of fodder sorghum M P Chari**

S. No.	Treatment combinations	Economics			
		Cost of cultivation	Gross return	Net return	B:C ratio
1.	Nitrogen 60 kg/ha + Spacing 35x15cm	27206	67397	40191	1.48
2.	Nitrogen 60 kg/ha + Spacing 45x15cm	26006	70466	44460	1.71
3.	Nitrogen 60 kg/ha + Spacing 55x15cm	24806	73260	48454	1.95
4.	Nitrogen 80 kg/ha + Spacing 35x15cm	27636	70070	42434	1.54
5.	Nitrogen 80 kg/ha + Spacing 45x15cm	26436	73766	47330	1.79
6.	Nitrogen 80 kg/ha + Spacing 55x15cm	25236	80234	54998	2.18
7.	Nitrogen 100 kg/ha + Spacing 35x15cm	28076	76714	48638	1.73
8.	Nitrogen 100 kg/ha + Spacing 45x15cm	26876	81153	54277	2.02
9.	Nitrogen 100 kg/ha + Spacing 55x15cm	25676	83809	58133	2.26
10.	Control	26436	69454	43018	1.63

\*Data Was Not Subjected to the Statistical Analysis



**Fig. 1. Field investigation**

### 3.2 Economic Analysis

#### 3.2.1 Gross returns

Observations regarding the economics of treatments are given in Table 2.

Highest gross return (83809.00 INR/ha) was obtained in treatment-9 (Nitrogen 100 kg/ha + Spacing 55x15cm) as compared to other treatments.

#### 3.2.2 Net returns

Net return (58133.00 INR /ha) was found to be highest in treatment-9 (Nitrogen 100 kg/ha + Spacing 55x15cm) as compared to other treatments.

#### 3.2.3 Benefit cost ration

Benefit Cost ratio (2.26) was found to be highest in treatment-9 with (Nitrogen 100 kg/ha + Spacing 55x15cm) as compared to other treatments.

### 4. CONCLUSION

It was concluded that with the application of nitrogen 100 kg/ha and along with the spacing 55x15cm (Treatment-9), has performs positively and improves growth and yield parameters. Higher green fodder yield and higher gross return, higher net return and benefit cost ratio were also recorded with the application of nitrogen 100 kg/ha and along with the spacing 55x15 cm (Treatment-9). These findings are

based on one season therefore; further trials may be required for further confirmation.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Hoffman PC, Esser NM, Bauman LM, Denzine SL, Engstrom M, Jonos HC. Effect of dietary protein on growth and N balance of Holstein heifer. *Journal of Dairy Science*. 2001;84: 843-847.
2. Pushpendra Singh, Sumeriya HK. Effect of nitrogen on yield, economics and quality of fodder sorghum genotypes. *Ann. Pl. Soil Res*. 2012;14:133-135.
3. Iqbal A, Ayub M, Zaman H, Ahmad R. Impact of nutrient management and legumes association on agro-qualitative traits of maize forage. *Pakistan. J. Bot*. 2006;38:1079- 1084.
4. Ayub M, Tanveer A, Ali S, Nadeem MA. Effect of different nitrogen levels and seed rates on growth, yield and quality of sorghum (*Sorghum bicolor* L.) fodder. *Indian Journal Agricultural Sciences*. 2002;72(11):648-656.
5. Saleem MF, Randhawa MS, Hussain S, Wahid MA, Anjum SA. Nitrogen management studies in autumn planted maize (*zea mays* L.) hybrids. *The Journal of Animal and Plant Sciences*. 2009; 19(3):140-143.
6. Lemerle D, Verbeek B, Diffy S. Influence of field pea (*Pisum sativum*) density on grain yield and competitiveness with annual rye grass (*Loliumrigidum*) in south-eastern Australia. *Australian Journal of Experimental Agriculture*. 2006;46:1465-1472.
7. Gomez KA, Gomez AA. Three or more factor experiment. (In:) *Statistical Procedure for Agricultural Research* 2nd ed. 1976;139 -141.
8. Mane Shivprasad, Rajesh Singh. Effect of planting geometry and different levels of nitrogen on growth, yield and quality of multicut fodder sorghum (*Sorghum bicolor* (L.) Monech), *Journal of Pharmacognosy and Phytochemistry*. 2017;6(4):896-899.
9. Dudhat MS, Savalia MG, Ramdevputra MV. Response of forage maize to nitrogen and phosphorus levels. *Forage Research*. 2004;30:34-35.

© 2023 Chand et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/98149>