



**Advances in Research**  
9(4): 1-8, 2017; Article no.AIR.33140  
ISSN: 2348-0394, NLM ID: 101666096



SCIENCE DOMAIN *international*  
[www.sciencedomain.org](http://www.sciencedomain.org)

# Physicochemical and Sensory Characteristics of Cows' Milk Butter Processed in Khartoum State, Sudan

Mohamed Osman Mohamed Abdalla<sup>1\*</sup>, Saniya Saleem Jabir Ahmed<sup>1</sup>  
and Siham Abdalla Rahamtalla<sup>1</sup>

<sup>1</sup>Department of Dairy Production, Faculty of Animal Production, University of Khartoum, Shambat, P.O.Box 32, Postal Code 13314, Khartoum North, Sudan.

## Authors' contributions

This work was carried out in collaboration between all authors. Author MOMA designed the study and performed the statistical analysis. Author SSJA wrote the protocol and the first draft of the manuscript. Author SAR managed the literature search. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/AIR/2017/33140

### Editor(s):

(1) Carmen Lizette del Toro Sanchez, Department of Food Research, University of Sonora, Mexico.

### Reviewers:

(1) Zhanibek Yessimbekov, Shakarim State University of Semey, Kazakhstan.

(2) Chen-Chin Chang, University of Kang Ning, Taiwan.

Complete Peer review History: <http://www.sciencedomain.org/review-history/18818>

Original Research Article

Received 31<sup>st</sup> March 2017  
Accepted 17<sup>th</sup> April 2017  
Published 27<sup>th</sup> April 2017

## ABSTRACT

**Aims:** This investigation was conducted to evaluate the physicochemical and sensory characteristics, fatty acid and cholesterol contents of butter processed in Khartoum State, Sudan.

**Methodology:** Butter was manufactured traditionally by farmers in Khartoum North (T1) and Omdurman (T2), in addition to butter manufactured commercially in a dairy plant (T3) and butter manufactured by researchers in the laboratory (T4). Physicochemical and sensory characteristics in addition to cholesterol level and free fatty acids profile were evaluated at 1, 15, 30, 45 and 60-day intervals for samples stored at 5.0±1.0°C, and at 1, 7, 14, 21 and 28-day intervals for samples stored at 25±2.0°C.

**Results:** The results showed high moisture, acidity and specific gravity in T1 (28.98%, 0.860% and 0.940 respectively), while fat, pH value and iodine index were high in T4 (82.5%, 4.53 and 29.00 mg I/100g, respectively) and solids-non-fat and melting point were high in T3 (2.01% and 28.32°C

\*Corresponding author: E-mail: [abutahany@outlook.com](mailto:abutahany@outlook.com), [abutahany@yahoo.com](mailto:abutahany@yahoo.com), [abutahany1960@yahoo.com](mailto:abutahany1960@yahoo.com);

respectively). The saponification index was high in T2 and T3 (200.00 and 201.00 mg KOH/g respectively). The fatty acids C6:0 – C18:0, C16:1, C18:1 – C18:3, C20:0 were detected in varying concentrations. The cholesterol content was high in butter of T4 (166.7 mg/100 g). Butter of T1 was best in flavour (8.43) and overall acceptability (9.00), while butter of T3 was best in texture and colour (8.86).

**Conclusion:** There were variations in physicochemical and sensory characteristics in addition to cholesterol level and fatty acids content of butter manufactured by different methods.

*Keywords: Butter; cholesterol; fatty acid; physicochemical; sensory.*

## 1. INTRODUCTION

Butter is defined as the product made exclusively from milk or cream or both, with or without addition of colouring material and containing not less than 80% by weight of milk fat and more than 400 fatty acids, of which 66% are saturated, 30% monounsaturated and 4% polyunsaturated [1]. Stage of lactation, diet, dietary supplementation, season of the year and diseases have an impact on the fatty acid composition of milk fat. Fermentation is one of the oldest food processing techniques in the world which has been adopted across generations and most of these techniques produce an inferior final product of low yield and stability and require high labour input. The availability of small scale processing techniques is the important factor at the traditional farmer level for manufacture of high quality dairy products with good characteristics [2,3].

Traditional butter which is locally called "*zebda beladi*" is one of the products resulting from spontaneous fermentation and rich in fat. Churning is the most important step for butter making, during which the oil-in-water emulsion is broken, leading to an aqueous phase separation and formation of water-in-oil emulsion [4]. Despite the application of preventive measures during its production and distribution, traditional butter has a limited shelf-life, and the deterioration (oxidation) may be usually due to oxidation of unsaturated glycerides resulting in increased level of free fatty acids. Most of the knowledge about the traditional fermentation has not been documented, thus, it is in danger of being lost since technologies are evolving and families are forgetting about the traditional food preservation practices [5]. The quality of stored butter is governed by such factors as cream ripening, the manufacturing process, good hygienic practice during processing and handling, the storage temperature and the type of animal feeding [6].

Lipid oxidation includes fatty acid oxidation which generates compounds that affect food quality and even nutrition and food safety. Oxidative rancidity or autoxidation cannot be stopped by lowering temperature of storage since it is a chemical reaction with low activation energy. Research concerning oxidative deterioration has been pursued for many years but it has been given a boost by recognition that such oxidation can cause damage to cell membranes and DNA which may be involved in aging process and cancer [7,8]. Butter is evaluated for its rich flavour; many of compounds that play a role in characteristics of flavour of butter have been identified although there is currently no chemical mixture that has been able to replace the aroma of butter [9]. Diacetyl, lactones, short chain fatty acids, lipid-derived aldehydes and dimethyl sulfide are the components responsible for butter flavour. Sensory analysis has been employed as a method to evaluate the acceptable butter flavour. Previous studies have relied on grading and product shelf life to compare products or evaluate quality of butter and margarine and spread products [10].

In the traditional method, raw milk is left at room temperature ( $25\pm 2^\circ\text{C}$ ) until coagulation takes place after 18 hr. During the gelation step, the product is called "*laben rayeb*", which is separated by churning into an aqueous fraction "*rob*" and fat fraction "*furssah*". The containers used for fermentation and churning are the "*si'in*" and the "*bukhssa*". The former is a leather bag from whole goat or sheep skin, while the latter is a large gourd with a lidded narrow mouth [11]. Due to the complexity steps to obtain butter products and for different processing methods used in manufacture of butter in Sudan, it is necessary to have accurate physicochemical evaluation and sensory characteristics that determine the quality and acceptability of the butter. This study was conducted to evaluate the physicochemical and sensory characteristics of butter produced traditionally and commercially in Khartoum State, Sudan.

## 2. MATERIALS AND METHODS

### 2.1 Butter Manufacture

#### 2.1.1 Butter traditionally manufactured by dairy farmers in Khartoum North and Omdurman (T1 and T2 respectively)

Raw milk was left at room temperature ( $25\pm 2^{\circ}\text{C}$ ) overnight to coagulate and convert into "*laben rayeb*". Next morning "*laben rayeb*" was churned in a skin bag from goat called "*siin*" until the coalescence of fat globules was formed indicating the conversion of milk into butter "*furssah*" and a by-product called "*rob*". Butter "*furssah*" was stored at  $4^{\circ}\text{C}$ , and analysis was carried out within 24 hr.

#### 2.1.2 Butter commercially manufactured in a dairy plant (T3)

Sweet cream obtained by cream separation of fresh milk (30% fat) was cooled at  $4^{\circ}\text{C}$  till used. The cream was pasteurized at  $82^{\circ}\text{C}$  for 30 min, cooled to  $10^{\circ}\text{C}$  and held at this temperature for two hours (sometimes overnight) till the butter was manufactured. The cream temperature was adjusted to  $11.6^{\circ}\text{C}$  and churning was started by filling the churn with  $\frac{2}{3}$  volume followed by addition of cold water, and churning was continued until granules were formed. Butter was washed with cold water for 5 min, repeating the washing twice, then the washed water was drained off and butter was wrapped, weighed, packed and stored at  $4^{\circ}\text{C}$ , and analysis was carried out within 24 hr.

#### 2.1.3 Butter manufactured by the investigators (T4)

Butter was manufactured in the dairy processing laboratory, Department of Dairy Production, Faculty of Animal Production, University of Khartoum. Raw milk was obtained from the University of Khartoum dairy farm. After fat determination, milk was warmed to  $40^{\circ}\text{C}$  and the cream was separated by hand separator. The cream obtained was analyzed for fat content and kept overnight in the refrigerator ( $4^{\circ}\text{C}$ ). Next morning the cream was churned at  $12^{\circ}\text{C}$  by hand churner till whipped cream became coarser and semi-solid butter granules were formed that rapidly increased in size and separated sharply from the liquid buttermilk. Butter was washed with cold water several times and the excess water was removed. Butter was filled in sterile

disposable polyethylene bags and stored at  $4^{\circ}\text{C}$ , and analysis was carried out within 24 hr.

### 2.2 Sample Preparation

To ensure uniform distribution of water, the samples were warmed in unopened airtight containers in a water bath at  $35^{\circ}\text{C}$ , agitated to facilitate melting and the test sample was thoroughly mixed to a homogeneous form.

### 2.3 Sample Analysis

Physicochemical (moisture, fat, solids-non-fat (SNF), pH, acidity, specific gravity, melting point, iodine index, saponification index, fatty acid composition, cholesterol content) and sensory characteristics (colour, flavour, texture, overall acceptability) were determined at the end of storage period of 60 days.

### 2.4 Physicochemical Composition

The fat, moisture, SNF and acidity were determined according to AOAC [12]. Melting point, iodine index, saponification index and specific gravity were determined according to FSSAI [13]. Fatty acids were determined by gas chromatography (Shimadzu 8A equipped with flame ionization detector) according to the method described by Radwan [14]. Cholesterol content was determined by an enzymatic method for the determination of total serum cholesterol [15].

### 2.5 Sensory Evaluation of Butter

Butter samples were subjected to descriptive sensory analysis using the 9-point hedonic scale [16], where 9 = excellent, 8 = very good, 7 = good, 6 = slightly good, 5 = fair, 4 = poor, 3 = slightly poor, 2 = very poor, 1 = extremely poor. Fourteen panelists were chosen to evaluate butter samples for colour, texture, flavour and overall acceptability.

### 2.6 Statistical Analyses

The data were subjected to descriptive statistical analysis using Statistical Analysis Systems (SAS, ver. 9). One-way randomized complete design was used for physicochemical analysis, fatty acid profile, cholesterol content and sensory analysis. Means were separated by Duncan multiple range test at  $P\leq 0.05$ .

### 3. RESULTS AND DISCUSSION

#### 3.1 Physicochemical Properties of Butter Samples

There was a significant ( $P < 0.05$ ) effect of treatment on the physicochemical properties of butter (Table 1). The highest fat content (82.50%) was found in butter manufactured by researchers in the laboratory (T4), while the lowest fat content (69.75%) was found in traditional farms in Khartoum North (T1). The content of fat in butter obtained from traditional farms in Khartoum north, Sudan is higher than those reported by Samet-Bali et al. [17] who found that fat content of traditional Tunisian butter was  $65.70 \pm 2.16\%$ . The moisture content ranged between 28.98% in T1 and 16.77% in T4. The moisture is higher in the traditional butter than the limits established for industrial butter. These findings are in agreement with previous studies [18,19], and lower than those reported by Kacem and Karam [20]. SNF content of butter was as high as  $2.01 \pm 0.01\%$  in T3 and as low as 0.90% in T4. The results of T1, T2 and T4 are lower than those reported by Makawi [21] who indicated that, the average solids-non-fat content was 2.04%, while the results of T3 are similar to Makawi [21]. Butter in T3 and T4 was less acidic (0.280%) than the other treatments (0.860% and 0.595% for T1 and T2 respectively). The results in this study are in agreement with the findings of Kacem and Karam [20] who reported the titratable acidity (TA) of butter to be 0.36 -0.90%. Meshref [22] reported a TA value of 0.04-0.55%

in cooking butter in Beni-Suef governorate, Egypt. In this study, the specific gravity of butter ranged between 0.900 in T2 and 0.940 in T1. The highest melting point was in butter from T3 (28.32°C), while the lowest value (27.44°C) was in T1. These values are in line with those reported by Asresie et al. [19] who reported values of  $11.0 \pm 0.100 - 42.9 \pm 1.00^\circ\text{C}$  for butter made from camel milk alone and blending it with goat milk. Significantly ( $P < 0.05$ ) higher iodine index was found in T4 (29.00), while the lower index was in T2 (26.25 mg I /100 g). The results of this study are lower than those obtained by Idoui et al. [18] who reported a value of 35.35 - 48.33 mg I/100 g in traditional Algerian butter from cow milk, and Idoui et al. [23] who reported a value of 37.17 - 85.95 mg I /100 g in Algerian traditional butter from goat milk. However, the results in this study are higher than those reported by Rady and Bader [24] who reported that the iodine index of cow's butter samples was 24.87-25.12 mg I /100 g, and that the value was not altered neither by gamma irradiation nor cold storage. The saponification index of butter was significantly ( $P < 0.01$ ) higher (200.00 mg KOH/g) in T2 and lower (192.90 mg KOH/g) in T1. The results are in accordance with those reported by Idoui et al. [18,23] for traditional Algerian butter from cows and goats respectively.

#### 3.2 Free Fatty acid Profile

Butter samples from all treatments were subjected to gas chromatography and lipids

**Table 1. Physicochemical characteristics of cow's milk butter manufactured in Khartoum state, Sudan**

Physicochemical characteristics	Treatment <sup>1</sup>				CV	SE	p
	T1	T2	T3	T4			
Fat (%)	69.75 <sup>c</sup>	70.72 <sup>c</sup>	79.80 <sup>b</sup>	82.50 <sup>a</sup>	0.17	0.0454	0.0409
Moisture (%)	28.98 <sup>a</sup>	28.20 <sup>a</sup>	18.19 <sup>b</sup>	16.77 <sup>c</sup>	1.37	0.0225	0.0112
Solids-non-fat (%)	1.27 <sup>b</sup>	1.07 <sup>c</sup>	2.01 <sup>a</sup>	0.90 <sup>d</sup>	8.94	0.0683	0.0234
Acidity (% lactic acid)	0.860 <sup>a</sup>	0.595 <sup>b</sup>	0.280 <sup>c</sup>	0.280 <sup>c</sup>	2.15	0.0002	0.0221
Specific gravity	0.940 <sup>a</sup>	0.900 <sup>c</sup>	0.910 <sup>b</sup>	0.903 <sup>c</sup>	1.04	0.0002	0.0195
pH value	4.36 <sup>b</sup>	4.30 <sup>c</sup>	4.33 <sup>c</sup>	4.53 <sup>a</sup>	0.22	0.0002	0.0432
Melting point (°C)	27.44 <sup>c</sup>	28.08 <sup>b</sup>	28.32 <sup>a</sup>	27.50 <sup>c</sup>	0.15	0.0258	0.0334
Iodine index (mg I/100g)	27.12 <sup>b</sup>	26.25 <sup>c</sup>	27.00 <sup>b</sup>	29.00 <sup>a</sup>	2.94	0.4715	0.0020
Saponification index (mgKOH/g)	192.90 <sup>c</sup>	200.00 <sup>a</sup>	198.00 <sup>b</sup>	201.00 <sup>a</sup>	0.41	0.4715	0.0290

Means in each row bearing similar superscripts are not significantly different ( $P > 0.05$ ), CV = Coefficient of variation (%), SE = Standard error of means, <sup>1</sup>Treatments 1, 2, 3 and 4 refer to butter manufactured in Khartoum North, Omdurman, the dairy plant and the laboratory respectively

resolved were determined to their constituent's fatty acids. The results showed that 12 fatty acids were identified; C<sub>6:0</sub>, C<sub>8:0</sub>, C<sub>10:0</sub>, C<sub>6:14</sub>, C<sub>14:0</sub>, C<sub>16:0</sub>, C<sub>16:1</sub>, C<sub>18:0</sub>, C<sub>18:1</sub>, C<sub>18:2</sub>, C<sub>18:3</sub> and C<sub>20:0</sub> (Table 2). Total saturated fatty acids (SFA%) are 55.81, 57.68, 56.28 and 56.03, while total mono- and poly-unsaturated fatty acids (MUFA+PUFA%) are 44.19, 42.31, 43.72 and 43.97 for T1, T2, T3 and T4 respectively. Important differences were found in fatty acids composition among butter samples, which contained a high percentage of saturated fatty acids (SFA), with the palmitic acid being the major (21.12-24.23%) followed by stearic acid (11.93-14.66%) and myristic acid (9.61-10.06%). These results are in line with those reported by Rady and Bader [24] who reported that, palmitic acid was the major SFA (22.81%) in butter followed by stearic acid (10.21%). Idoui et al. [18] reported that traditional cows' butter contained a high percentage of SFA and palmitic acid was the major SFA (24.33-36.95%) followed by myristic acid (18.49-27.35%) and stearic acid (7.68-14.05%). The predominance of palmitic acid in butter milk fat was also reported in previous studies [18,23]. In a study conducted by Ito et al. [25], it was found that heated butter contained 27.06% palmitic. Oleic acid was the predominant mono-unsaturated fatty acid (34.16-35.41%). Total saturated fatty acids in this study were higher than those reported by Rady and Bader [24], while total unsaturated FA were lower. Butyric acid (C<sub>4:0</sub>) is the most prevalent short-chain fatty acid and often identified with

butter due to its characteristics impact on flavour. In this study, butyric acid was not identified, probably due to its volatility. Some of the fatty acids such as caproic, caprylic, capric, lauric, palmitoleic, linoleic, linolenic and arachidic were present in samples from all treatments.

### 3.3 Cholesterol Content

Cholesterol content was significantly affected by the source of butter samples, being higher (166.7 mg/100g) in T4 and lower (136.7 mg/100g) in T2 (Table 3). Cholesterol is indispensable for life, it serves to build and maintain the membranes that surround the cells, and it is necessary for the synthesis of some hormones especially sexual ones. It also contributes in producing biliary acid that plays a role in digestion of fat, in addition to contributing in biosynthesis of vitamin D [26]. Milk and milk products play an important part in healthy diet as they contribute to intake of essential nutrients and protein of a high nutritional quality. However, due to high content of cholesterol raising saturated fatty acids (SFA) in milk fat, decrease in the intake of fat-rich dairy products is recommended, although there is no doubt that mixture of long-chain saturated fatty acids in milk increases plasma cholesterol [27]. Rady and Badr [24] reported that cholesterol is the only sterol compound showing a slight decrease upon the application of accident doses  $\gamma$ -rays which may be attributed to partial slight oxidation.

**Table 2. Fatty acid profile (%) of cow's milk butter manufactured in Khartoum state, Sudan**

Fatty acid	Symbol	Treatment <sup>1</sup>				CV	SE	p
		T1	T2	T3	T4			
Caproic	C6:0	1.355 <sup>b</sup>	1.627 <sup>a</sup>	1.305 <sup>b</sup>	1.500 <sup>ab</sup>	8.16	0.0683	0.0212
Caprylic	C8:0	1.882 <sup>d</sup>	1.996 <sup>c</sup>	1.677 <sup>e</sup>	2.250 <sup>a</sup>	1.96	0.0182	0.0442
Capric	C10:0	4.378 <sup>bc</sup>	4.368 <sup>bc</sup>	3.914 <sup>cd</sup>	3.666 <sup>d</sup>	5.68	0.1494	0.0409
Lauric	C12:0	3.116 <sup>b</sup>	3.308 <sup>b</sup>	2.982 <sup>b</sup>	3.000 <sup>b</sup>	12.55	0.2588	0.0329
Myristic	C14:0	9.61 <sup>a</sup>	9.59 <sup>a</sup>	10.06 <sup>a</sup>	9.50 <sup>a</sup>	3.27	0.1713	0.0449
Palmitic	C16:0	21.74 <sup>bc</sup>	22.05 <sup>b</sup>	24.23 <sup>a</sup>	21.12 <sup>cd</sup>	2.36	0.2834	0.0119
Palmitoleic	C16:1	3.564 <sup>b</sup>	3.229 <sup>c</sup>	2.795 <sup>d</sup>	3.749 <sup>b</sup>	4.44	0.0948	0.0129
Stearic	C18:0	13.53 <sup>b</sup>	14.60 <sup>b</sup>	11.93 <sup>c</sup>	14.66 <sup>b</sup>	5.24	0.4367	0.0385
Oleic	C18:1	34.49 <sup>ab</sup>	34.38 <sup>ab</sup>	35.41 <sup>a</sup>	34.16 <sup>ab</sup>	2.11	0.4054	0.0299
Linoleic	C18:2	5.755 <sup>b</sup>	4.299 <sup>d</sup>	5.143 <sup>bc</sup>	5.557 <sup>b</sup>	6.08	0.2244	0.0400
Linolenic	C18:3	0.3727 <sup>c</sup>	0.4040 <sup>c</sup>	0.3730 <sup>c</sup>	0.5000 <sup>b</sup>	5.56	0.0182	0.0291
Arachidic	C20:0	0.1893 <sup>d</sup>	0.1643 <sup>f</sup>	0.1860 <sup>e</sup>	0.3330 <sup>b</sup>	4.25	0.0001	0.0333
Total saturated	-	55.81 <sup>b</sup>	57.68 <sup>a</sup>	56.28 <sup>b</sup>	56.03 <sup>b</sup>	0.93	0.3028	0.0400
Total unsaturated	-	44.19 <sup>b</sup>	42.31 <sup>c</sup>	43.72 <sup>b</sup>	43.97 <sup>b</sup>	1.21	0.3055	0.0126

Means in each column (capital letter) and row (small letter) bearing similar superscripts, are not significantly, different ( $P>0.05$ ). CV = Coefficient of variation (%), SE = Standard error of means, <sup>1</sup>Treatments 1, 2, 3 and 4 refer to butter manufactured in Khartoum North, Omdurman, the dairy plant and the laboratory respectively.

**Table 3. Cholesterol content of cow's milk butter manufactured in Khartoum state, Sudan**

Treatment1	Cholesterol content (mg/100g)
T1	146.7c
T2	136.7e
T3	142.0d
T4	166.7b
CV (%)	6.97
SE	0.3978
<i>p</i>	0.0081

Means in column bearing similar superscripts are not significantly different ( $P>0.05$ ). CV = Coefficient of variation, SE = Standard error of means, <sup>1</sup>Treatments 1, 2, 3 and 4 refer to butter manufactured in Khartoum North, Omdurman, the dairy plant and the laboratory respectively

**Table 4. Sensory evaluation of cow's milk butter manufactured in Khartoum state, Sudan**

Treatment1	Sensory attributes			
	Flavour	Texture	Colour	Overall acceptability
T1	8.43 <sup>a</sup>	8.43 <sup>b</sup>	6.57 <sup>d</sup>	9.00 <sup>a</sup>
T2	7.85 <sup>b</sup>	8.14 <sup>c</sup>	6.28 <sup>e</sup>	8.43 <sup>c</sup>
T3	8.14 <sup>ab</sup>	8.86 <sup>a</sup>	8.86 <sup>a</sup>	8.71 <sup>b</sup>
T4	7.43 <sup>c</sup>	6.71 <sup>d</sup>	8.57 <sup>b</sup>	8.71 <sup>b</sup>
CV (%)	5.49	4.87	5.08	5.34
SE	0.0831	0.0532	0.0429	0.0576
<i>p</i>	0.0236	0.0443	0.0376	0.0165

Means in each column bearing similar superscripts are not significantly different ( $P>0.05$ ). \* =  $P<0.05$ , CV = Coefficient of variation, SE = Standard error of means, <sup>1</sup>Treatments 1, 2, 3 and 4 refer to butter manufactured in Khartoum North, Omdurman, the dairy plant and the laboratory respectively

### 3.4 Sensory Evaluation

Flavour of butter samples in T1 significantly scored the highest (8.43), while samples in T4 scored the lowest (7.43). However, the samples in T3 had the best texture (8.86) followed by samples in T1, T2 and T4 respectively. The colour of samples in T3 was the best (8.86) and the colour of samples in T2 was the worst (6.28). Overall, butter samples in T1 were highly acceptable (9.0), while samples in T2 were the least acceptable (8.43) (Table 4 above). In this study, the greatest difference was observed for colour, ranging from 6.28 to 8.86. The butter samples after storage were highly accepted by descriptive analysis and this is in agreement with Krause et al. [28] who reported that, for optimum quality, butter quainter should be refrigerated for at least 6 months; but when frozen at -20 °C, stick can be stored up to 2 years, and for bulk butter in refrigerated containers, flavour quality is maintained for at least 9 months. Fernandes et al. [29] reported that the colour of butter is not only determined by the level of carotene in the milk fat, but also by the size distribution of water droplets, and the finer the droplets, the greater the scattering of the light, and therefore the lighter the colour of the butter. Several studies

revealed that, although texture changes with milk fatty acid composition, the effect on flavour is not significant [30,31]. Rady and Bader [24] reported that, mean scores of colour and appearance of cow's butter samples was found as well as flavour (including aroma, taste and texture). Post-treatment and upon refrigeration storage, it is obvious that gamma irradiation had no adverse effect on the sensory quality attributes of butter [24].

### 4. CONCLUSION

From the results obtained, it could be concluded that, the general prerequisites for high quality butter are influenced by the different methods of manufacturing cow's milk butter in Sudan. The desirable flavour and overall acceptability in terms of sensory properties and high content of unsaturated fatty acids is the major feature of butter manufacture by the traditional methods. This could highlight the role of this section to provide a good quality butter for consumers in Sudan.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. USDA. United States Standards for Grades of Butter. Section 58-2621. Available:[http://www.ams.usda.gov/AMSV1\\_0/getfile?dDocName=STELDV3004470](http://www.ams.usda.gov/AMSV1_0/getfile?dDocName=STELDV3004470) (1989).
2. O'Mahony F, Peters JK. Options for smallholder in Sub-Saharan Africa. ILCA Bulletin. Addis Ababa, Ethiopia: International Livestock Research Institute (ILRI). 1987;27:2-17.
3. Kebede A, Viljoen BC, Gadaga TH, Narvhus JA, Lourens-Hattingh A. The effect of container type on the growth of yeast and lactic acid bacteria during production of Sethemi, South African spontaneously fermented milk. Food Research International. 2007;40(1):33-38.
4. Rousseau D. Fat crystals and emulsion stability- A review. Food Research International. 2000;33(1):3-14.
5. Muir DD. The shelf-life of dairy products: 3. Factors influencing intermediate and long life dairy products. International Journal of Dairy Technology. 1996;49(3): 67-72.
6. Fearon AM, Mayne CS, Charlton CT. Effect of naked oats in the cow's diet on the oxidative stability of the milk fat. Journal of the Science of Food and Agriculture. 1998;76(4):546-552.
7. Moller P, Wallin H. Adduct formation, mutagenesis and nucleotide excision repair of DNA damage produced by reactive oxygen species and lipid peroxidation product. Mutation Research/ Reviews in Mutation Research. 1998; 410(3):271-290.
8. Tonfa M, Sevastita M, Sonia S, Elena M, Constanta M. Researches regarding the quantitative modifications of hop pellets bitter acids during the storage. Bulletin USAMV-CN. 2006;62:365-370.
9. Peterson DG, Reineccius GA. Characterization of volatile compounds that constitute fresh sweet cream butter aroma. Flavour and Fragrance Journal. 2003;18(3):215-220.
10. Thompson JL, Darke MA, Lopetcharat K, Yates MD. Preference mapping of commercial chocolate milks. Journal of Food Science. 2004;69(9):S406-S413.
11. Abdelgadir WS, Tagelsir KA, Dirar HA. The traditional fermented milk products of the Sudan: Review. International Journal of Food Microbiology. 1998;44:1-13.
12. AOAC. Official methods of analysis, 15th edition. Washington, D.C.: The Association Official Analytical Chemists, USA; 2000.
13. FSSAI. Manual of methods of analysis of foods (oils and fats). In: Food Safety and Standards Authority of India. New Delhi: Ministry of Health and Family Welfare, Government of India; 2012.
14. Radwan SS. Coupling of two dimensional thin-layer chromatography with gas chromatography for the quantitative analysis of lipid classes and their constituent fatty acids. Journal of Chromatography Science. 1978;16:538-542.
15. Allain CC, Poon LS, Chan CS. Enzymatic determination of total serum cholesterol. Clinical Chemistry. 1974;20(4):470-475.
16. Lim J. Hedonic scaling: A review of methods and theory. Food Quality and Preference. 2011;22:733-747.
17. Samet-Bali O, Ayadi MA, Attia H. Traditional Tunisian butter, physicochemical and microbial characteristics and storage stability of the oil fraction. Food Science and Technology. 2009;42:899-905.
18. Idoui T, Benhamada N, Leghouchi E. Microbial quality, physicochemical characteristics and fatty acid composition of traditional butter produced from cow's milk in east Algeria. Grasas y Aceites. 2010;61(3):232-236.
19. Asresie A, Seifu E, Kurtu MY. Physicochemical properties of butter made from camel milk alone and blending it with goat milk. World Journal of Animal Science Research. 2013;1(1):1-8.
20. Kacem M, Karam NE. Physicochemical and microbiological study of "shmen", a traditional butter made from camel milk in the Sahara (Algeria): Isolation and identification of lactic acid bacteria and yeasts. Grasas y Aceites. 2006;57(2):198-204.
21. Makawi EA. Chemical composition, keeping quality and yield of butter at Kuku dairy plant. M.Sc. Thesis Khartoum, University of Khartoum, Sudan; 1992.
22. Meshref SMA. Microbiological quality and safety of cooking butter in Beni-Suef governorate, Egypt. African Health Science. 2010;10(2):193-198.
23. Idoui T, Rechak H, Zabayou N. Microbial quality, physicochemical characteristics and fatty acid composition of traditional butter produced from cow's milk. Annals.

- Food Science and Technology. 2013; 14(1):108-114.
24. Rady AH, Badr HM. Keeping the quality cows' butter by  $\gamma$ -irradiation. *Grasas y Aceites*. 2003;54:410-418.
25. Ito N, Wada S, Yamanaka Y, Takagaki H, Nakamura H. Identification of novel decanoic acid in heated better. *Bioscience, Biotechnology and Biochemistry*. 2005;69: 2416-2420.
26. Cerin E, Leslie E, Sugiyama T, Owen N. Association of multiple physical activity domains with mental well-being. *Mental Health and Physical Activity*. 2005;6(3): 363-376.
27. Wood R, Kubena K, O'Brien B, Tseng S, Martin G. Effect of butter, mono and poly unsaturated fatty acids-enriched butter, trans fatty acids margarine, and zero trans fatty acids margarine on serum lipids and lipoproteins in healthy men. *Journal of Lipid Research*. 1993;34:1-11.
28. Krause AJ, Lopetcharat K, Drake MA. Identification of the characteristics that drive consumer liking of the butter. *Journal of Dairy Science*. 2007;90:2091-2102.
29. Fernandes SAA, Ramos DS, Ramos EM, Veras DVS, Pinheiro FR, Lag Neto S, Requião LA, Amui TS, Carneiro JCS, Matarazzo SV. Sensory evaluation of buffalo butter. *Italian Journal of Animal Science*. 2010;6:1140-1142.
30. Lin MP, Sims CA, Staples GR, O'keefe SF. Flavor quality and texture modified fatty acid high monoene, low saturate butter. *Food Research International*. 1996;29(3-4):367-371.
31. Bobe G, Hammond EG, Freeman AE, Lindberg GL. Texture of butter from cows with different milk fatty acid compositions. *Journal of Dairy Science*. 2003;86(10): 3122-3127.

© 2017 Abdalla et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://sciencedomain.org/review-history/18818>