



Evaluation of Silicon Nutrition on Yield, Growth Attributes and Available Nutrient Status of Paddy in Coastal Zone of Odisha

Aliva Das ^a, Prasanna Kumar Samant ^a, Gayatri Sahu ^{b*} and Gour Hari Santra ^b

^a Department of Soil Science and Agricultural Chemistry, Odisha University of Agriculture and Technology, Bhubaneswar, Odisha- 751003, India.

^b Department of Soil Science and Agricultural Chemistry, Institute of Agricultural Sciences, Siksha 'O' Anusandhan Deemed to be University, Bhubaneswar, Odisha- 751030, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2021/v33i2330742

Editor(s):

(1) Prof. RusuTeodor, University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, Romania.

Reviewers:

(1) Chuck Chuan NG, Xiamen University Malaysia, Malaysia.

(2) Dionisio S. Bucao, Mariano Marcos State University, Philippines.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:
<https://www.sdiarticle5.com/review-history/77740>

Original Research Article

Received 11 September 2021
Accepted 23 November 2021
Published 29 November 2021

ABSTRACT

A field experiment was conducted in the central farm, Regional Research and Technology Transfer Station, Coastal Zone OUAT, Bhubaneswar, Odisha, India during *kharif* 2020 using *cv-Lalat* variety of Rice. The experiment was laid out in randomized block design (RBD) with ten treatments and three replications. In this experiment BOF (Basic Oxygen Furnace) slag was used as a source of silica for application to rice crop. 200, 300 and 400 kg SiO₂/ha was applied in combination with 50% STD (Soil Test Dose) and 75% STD to assess the efficacy of Silica with reduction in fertilizer dose by 50% and 25%. It was shown that increase in number of effective tillers (7.8) per plant and length of the panicle (25.8 cm), but no significant increase was marked in case of 1000 grain weight (gm.) over control. In case of grain yield, straw yield and harvest index, 100% STD (T₄) registered maximum (36.9q/ha) grain yield which was 29.4% more than that of control. Of course, grain yield in case of T₉ i.e., 75% STD+300 kg SiO₂/ha was at par 36.2q/ha indicated efficacy of silica application by reducing fertilizer dose. Harvest index was not affected by Si treatments. However, maximum HI (0.478) was observed in T₉. Available N, P, K and S status in post-harvest soil was increased due to application of silica along with fertilizer over control indicated better availability of

*Corresponding author: E-mail: gayatrisahu@soa.ac.in, gayatrickv@gmail.com;

nutrients which plays a vital role in increasing production and productivity of rice. The study revealed that the DTPA extractable Fe and Mn content decreased, and Cu and Zn content increased with increased application of silica. Though 100% STD (T_4) was found to be very effective as compared to other treatments but 75% STD + 300kgSiO₂/ha was also equally effective so far as yield, growth attributing characters, available nutrient status of paddy.

Keywords: Rice; silicon; growth; yield; nutrient status.

1. INTRODUCTION

Silicon (Si) is graded as the second-most abundant element (after oxygen) in the earth's crust with nearly 29% mean content [1]. The predominant form of Si in soil solution is monosilicic acid (H₄SiO₄) [2]. Iron and Al-oxides of soil have the capacity to adsorb a significant amount of Si on their surfaces. The potential of Si in improving crop yield has been established in many studies, especially under abiotic and biotic stress conditions like drought, heavy metals, salinity, and pathogens [3, 4, 5, 6]. Si is known for its role in alleviating the negative stress effects on many plant species. Rice (*Oryza sativa* L.) is a high Si accumulating plant [2]. Besides rice yield increase, Si has many fold advantages of increasing nutrient availability (N, P, K, Ca, Mg, S and Zn) [7], decreasing nutrient toxicity (Fe, Mn, and Al) and minimizing biotic and abiotic stress in plants [8]. Hence, the application of Si to soil or plant is practically useful in laterite derived rice soils, not only to increase yield but also to alleviate the Fe toxicity problems. In general farmers export silicon from field by removing straw residues with the harvest and the exogenous application of silicon in rice is overlooked. Therefore, a continued supply of Silicon would be required predominantly for the healthy and productive development of plant during all growth stages [9]. The use of silica fertilizers in the form either soluble silicates or of calcium silicate slag is still very restricted. Hence, effective practical means of application and affordable sources of Si are needed. With this background, the present study was undertaken to find out a suitable dose of SiO₂ for increasing growth and yield attributing character in rice by alleviating the toxicity of Fe, Mn, and Al in acidic soils.

2. MATERIALS AND METHODS

The experiment was conducted in Central Farm, Regional Research and Technology Transfer Station, O.U.A.T, Bhubaneswar between latitude 20.2961° N and longitude 85.8245° E and an average altitude of 45 m (148 ft) above sea level.

The experiment was carried out in a Randomised Block Design (RBD) with ten treatments replicated thrice. The treatments were T1- Control, T2- 50% STD (Soil Test Dose), T3- 75% STD, T4- 100% STD, T5- 50% STD+200 kg SiO₂/ha, T6- 50% STD+300 kg SiO₂/ha, T7- 50% STD+400 kg SiO₂/ha, T8- 75% STD+200 kg SiO₂/ha, T9- 75% STD+300kg SiO₂/ha, T10- 75% STD+400 kg SiO₂/ha. The soil of the experimental site was sandy, acidic reaction (pH 5.67), EC (0.16dS/m), O.C(0.41%), Av. N(178kg/ha), Av. P(13.5kg/ha), Av. K (83kg/ha), Av. S (6.8kg/ha), DTPA extractable Fe(78.8ppm), Mn(1.32ppm), Cu (0.21ppm) and Zn(0.35ppm). In the experiment BOF (Basic Oxygen Furnace) slag a byproduct of Jindal Steel Pvt. Ltd. was used as a source of silica for application to rice crop. 200, 300 and 400 kg SiO₂/ha was applied in combination with 50% STD and 75% STD to assess the effect of Silica with reduction in fertilizer dose by 25% and 50%. Rice cv. Lalat was selected for the research purpose since it suits for the medium land situation. The nutrients were applied in form of Urea, Diammonium phosphate (DAP) and Muriate of Potash (MOP) and Silicate Slag as per the experimental plan. The requirement of nitrogen by the crop was supplemented in the form of DAP and Urea in three splits viz, 25% basal, 50% at active tillering stage and finally 25% at panicle initiation stage. Required amount of phosphorus were applied as 100% basal in each plot. Total amount of potassium in form of Muriate of Potash (MOP) was applied in two splits i.e., 50% at basal and 50% at panicle initiation stage. Soil application of Silicate Slag was made @ 200, 300 and 400 kg SiO₂/ha at the time of transplanting as per the treatment imposed. Soil samples were analysed for available nutrient status before and after the harvest of the crop and expressed as kg ha⁻¹. Biometric observations i.e., growth and yield attributing parameters like plant height, number of tillers and panicles per hill, panicle length, grains per panicle, 1000 grain weight etc. were recorded at different growth stage and at harvest. The results obtained were statistically analysed using statistical analysis software (SAS) i.e., DOS-based SPSS version 10.0.

3. RESULTS AND DISCUSSION

3.1 Effect of Silica on Growth Attributing Characters of Paddy

The experimental data presented in Table 1 indicates influence of Silica on plant growth attributing characters like number of effective tillers per plant, length of the panicle and 1000 grain weight.

Number of effective tillers per plant varied from 5.5 to 8.4 in which lowest number of tillers was observed in T₁ i.e., Absolute control (No Fertilizer) and significantly maximum number of tillers was observed in T₄ i.e., 100% STD. Number of effective tillers increased as the fertilizer dose increased. In case of application of silica in combination with 50 and 75% STD, maximum number of tillers was obtained due to application of 300kg SiO₂/ha which was reduced with increased application of silica. 100% STD was superior to all other treatments but 75% STD+300 kg SiO₂/ha was also equally effective when all the characters taken together. Hence, by reducing the fertilizer dose by 25% and applying 300kg SiO₂/ha we can achieve the desired quality. In our study, the increase in total number of effective tillers per plant might be attributed to increased availability of phosphorus and other beneficial effect of silicon on growth of paddy. These results are in confirmation with those reported by those who also reported the beneficial role of Si fertilizer in increasing number of effective tillers per plant [10, 11, 12, 13, 14, 15].

Length of the panicle was varied from 22.5 cm to 25.8 cm. with lowest in T₁ (Absolute control). 100% STD (T₄) and 75% STD + 300 kg SiO₂/ha (T₉) registered significantly maximum panicle length which is 14.6% more than that of control. There was gradual increase in panicle length as the fertilizer dose increased. Application of 300kg SiO₂/ha registered maximum panicle length when compared with 200 and 400kg SiO₂/ha irrespective of integration of fertilizer dose. Of course, there was no significant difference between 300 and 400 kg SiO₂/ha so far as length of the panicle is concerned. The study shows an increase in length of panicle which is also in confirmation with some experiments [13, 16].

1000 grain wt. varied from 21.8gm to 23.1 gm. Application of silica. Increased fertilizer dose did not have any significant effect on 1000 grain wt. It was lowest in absolute control i.e., 21.8 gm and highest in 100% STD i.e., 23.1 gm which was at par with 75% STD + 300kg SiO₂/ha. There was no significant difference between the treatments. Application of Silica which contradicts the findings of some research works, which shows significantly increased 1000-grain weight with application of Si fertilizer [17, 13, 18, 15].

3.2 Effect of Silica on pH, Electrical Conductivity and Organic Carbon Content of Post-Harvest Soil

The experimental data presented in Table 2 indicates effect of Silica on pH, Electrical Conductivity and Organic Carbon in post-harvest soil. The study shows increase in pH, Electrical Conductivity (EC) and Organic Carbon (OC) in post-harvest soil due to application of Silica.

Table 1. Effect of Silica on number of effective tillers per plant, length of the panicle and 1000 grain weight

Treatment details	Number of effective tillers per plant	Length of the panicle (cm)	1000 grain wt.(gm)
T ₁ = Absolute Control	5.5	22.5	21.8
T ₂ = 50% STD	6.1	23.2	22.1
T ₃ = 75% STD	6.3	24.1	22.2
T ₄ = 100% STD	8.4	25.8	23.1
T ₅ = T ₂ + 200kg SiO ₂ /ha	6.5	24.5	22.4
T ₆ = T ₂ + 300kg SiO ₂ /ha	7.1	25.0	22.7
T ₇ = T ₂ + 400kg SiO ₂ /ha	6.6	24.7	22.4
T ₈ = T ₃ + 200kg SiO ₂ /ha	6.9	24.9	22.5
T ₉ = T ₃ + 300kg SiO ₂ /ha	7.4	25.8	23.0
T ₁₀ = T ₃ + 400kg SiO ₂ /ha	7.3	25.4	22.8
SE (M)	0.18	0.26	-
CD (0.05)	0.55	0.78	NS

[SE- Standard Error, CD- Critical Difference, NS- Non-Significant]

It was observed that that there was a variation in pH value so far as different treatments are concerned. pH value decreased with T₂ (50% STD) over T₁ but increased with further application of fertilizer i.e., T₃ (75% STD) and T₄ (100% STD) over control. pH value was recorded maximum (5.98) in T₁₀. Application of silicon in different doses along with T₂ (50% STD) and T₃ (75% STD) shows increase in pH value. However, this increase was not significant. This increase in soil pH could be attributed to the fact that silicate materials can increase soil reaction and help in correcting soil acidity by neutralizing exchangeable Fe, Al and Mn and other toxic elements [19]. These results were also in line with that reported by some scientists [20, 21, 22, 9].

Variation in Electrical Conductivity ranged from 0.118 dS/m to 0.148 dS/m. Electrical Conductivity was maximum in T₁₀ (75% STD+400kg SiO₂/ha) and minimum in control (0.118 dS/m). There was gradual increase in Electrical Conductivity of post-harvest soil as the fertilizer dose increased. However, the increase was not significant. Application of silica in different doses along with T₂ (50% STD) or T₃ (75% STD) showed an increasing trend. The silicon application in soil resulted in increase in soil Electrical Conductivity (EC) after the experiment. This might be attributed to submergence, increase in solubility of salts present in the soil and due to the dissolution of silicon fertilizers.

There was a variation ranging from 0.31 % to 0.48 % of OC in post-harvest soil so far as different treatments are concerned. Maximum OC content (0.48%) in soil was recorded in T₄ (100% STD) and T₇ (50% STD+ 400kg SiO₂/ha)

as compared to lowest in absolute control T₁. This increase in OC content in soil was 55% more over control. With increase in fertilizer doses, the percentage of OC was also increased. All the treatments registered significantly higher OC content over Control. In case of application of silica, in combination with 50% and 75% STD, the percentage of OC increases but the increase was not significant. Application of 400 kg SiO₂/ha registered maximum organic carbon content in soil when compared with 200 and 300 kg SiO₂/ha irrespective of integration of fertilizer dose. The increase in soil organic carbon (OC) was due to the reason that organic materials had direct impact on mineralization rate that increases soil organic carbon directly [23].

3.3 Effect of Silica on Available Soil Nutrients in Post-Harvest Soil

The experimental data presented in Table 3 indicates effect of Silica on available soil nutrients in post-harvest soils like Nitrogen (N), Phosphorous (P), Potash (K) and Sulphur (S). Analysis of initial soil sample indicated low available N (178 kg/ha), P (13.5kg/ha), K (83kg/ha) and S (6.8kg/ha) status which when compared with post-harvest soil analysis found a little increase due to application of Silica. Maximum available N(201.7kg/ha), P(14.5kg/ha), K (92.2 kg/ha) and S (16.2 kg/ha) status was observed in 75% STD+ 300kg SiO₂/ha which enlighten us about the beneficial effect of application of silica to soil when compared with control. It was stated that beside yield enhancement in rice, silicon also has many fold advantages of increasing availability of major nutrients and also alleviating iron toxicity problems in soil [24]. These results are confirmative with some findings [25,26, 27, 9].

Table 2. Effect of Silica on soil pH, Electrical Conductivity and Organic Carbon content of post-harvest soil sample

Treatment details	pH	E.C(dS/m)	O.C (%)
T ₁ = Absolute Control	5.41	0.118	0.31
T ₂ = 50% STD	5.20	0.121	0.42
T ₃ = 75% STD	5.48	0.130	0.45
T ₄ = 100% STD	5.64	0.135	0.48
T ₅ = T ₂ + 200kg SiO ₂ /ha	5.71	0.120	0.41
T ₆ = T ₂ + 300kg SiO ₂ /ha	5.85	0.124	0.44
T ₇ = T ₂ + 400kg SiO ₂ /ha	5.91	0.136	0.48
T ₈ = T ₃ + 200kg SiO ₂ /ha	5.89	0.142	0.42
T ₉ = T ₃ + 300kg SiO ₂ /ha	5.90	0.146	0.43
T ₁₀ = T ₃ + 400kg SiO ₂ /ha	5.98	0.148	0.46
SE (M)	0.13	0.006	0.03
CD (0.05)	0.39	0.018	0.09

[SE- Standard Error, CD- Critical Difference]

Table 3. Effect of Silica on available soil nutrients in post-harvest soil

Treatment details	Available Nutrients (Kg/ha)			
	N	P	K	S
T ₁ = Absolute Control	170.5	9.2	76.1	6.5
T ₂ = 50% STD	183.5	11.9	85.2	7.6
T ₃ = 75% STD	191.3	12.5	85.5	8.8
T ₄ = 100% STD	199.7	14.2	88.5	12.2
T ₅ = T ₂ + 200kg SiO ₂ /ha	184.5	13.2	84.6	10.4
T ₆ = T ₂ + 300kg SiO ₂ /ha	185.7	14.0	88.0	11.6
T ₇ = T ₂ + 400kg SiO ₂ /ha	186.2	13.5	86.2	12.8
T ₈ = T ₃ + 200kg SiO ₂ /ha	195.7	12.9	88.4	14.1
T ₉ = T ₃ + 300kg SiO ₂ /ha	201.7	14.5	92.2	16.2
T ₁₀ = T ₃ + 400kg SiO ₂ /ha	199.2	14.2	87.2	16.4
SE (M)	5.7	1.0	2.5	1.1
CD (0.05)	17.2	2.9	7.5	3.5

[SE- Standard Error, CD- Critical Difference]

Available N status ranged from 170.5 kg/ha to 201.7 kg/ha with lowest in T₁ i.e., Absolute control (No Fertilizer) and significantly maximum in T₉ i.e., 75% STD+ 300kg SiO₂/ha. Increased trend in available N status was observed as the fertilizer dose increased. All the treatments registered significantly higher available Nitrogen over Control except T₂ (50% STD). In case of application of silica in combination with 50 % STD, it was observed that available N status was increased with increased application of silica, and it was found maximum due to application of 400kg SiO₂/ha, but the increase was not significant. However, in case of application of silica in combination with 75 % STD, available N status was maximum (201.7 kg/ha) due to application of 300kg SiO₂/ha which was reduced with further application of silica. The study shows an increased dose of Si have its potential to raise the soil-available N and N-use efficiency and plant gained maximum benefits of ample N availability [12]. It was also found that application of Si enhanced the use of applied N to rice [28, 29]. This was in conformity with findings that Si application has the potential in enhancing the availability of N in soil [30, 12].

Available P status varied from 9.2 kg/ha to 14.5 kg/ha with a lowest value observed in T₁ i.e., Absolute control (No Fertilizer) and significantly maximum available P (14.5kg/ha) was observed in T₉ i.e., T₃+ 300kg SiO₂/ha which was 58 % higher over Control. T₄ (100% STD) and T₁₀ (75% STD+400kg SiO₂/ha) were at par with T₉. Available P status increased as the fertilizer dose increased. All the treatments registered significantly higher available P status over Control except T₂ (50% STD). In case of application of silica in combination with 50 % and

75 % STD, maximum available P (14.5kg/ha) was obtained due to application of 300kg SiO₂/ha which was reduced with further increased application of silica. It was also reported that better response of rice crops with respect to P nutrition might be due to increase in root growth and enhanced soil P availability with Si application, the decreased in P-retention capacity of soil, and the increased solubility of P, leading to increased efficiency of phosphatic fertilizers [31]. It was also reported an enhanced availability of P with the addition of Si fertilizer [12, 15].

Available Potassium status was varied from 76.1 kg/ha to 92.2 kg/ha. Lowest available K status was observed in T₁ i.e., Absolute control (No Fertilizer) and T₉ i.e., T₃+ 300kg SiO₂/ha registered significantly maximum available K which was 21 % higher over Control. Available K status was increased as the fertilizer dose increased. All the treatments registered significantly higher available K status over Control T₁. In case of application of silica in combination with 50 % and 75 % STD, maximum available K was obtained due to application of 300kg SiO₂/ha which was reduced with increased application of silica. Furthermore, it should be noted that Si application can alleviate the K-deficiency-induced growth inhibition by improving plant-water status via enhancement of stomatal conductance and transpiration rates [32]. The production of hydrogen ions during reduction of Fe and Al might have helped in the release of K from the exchange sites or from the fixed pool to the soil solution [9].

Available S status varied from 6.5 kg/ha to 16.4 kg/ha with lowest in T₁ i.e., Absolute control and

maximum in T₁₀ i.e., T₃+ 400kg SiO₂/ha. As the fertilizer dose increased, the available S status was also increased but no significant difference was marked up to 75% STD. In case of application of silica in combination with 50 % and 75 % STD, available S status was increased with increased application of silica which was significantly higher over Control. Application of 400kg SiO₂/ha registered maximum S status when compared with 200 and 300kg SiO₂/ha irrespective of integration of fertilizer dose.

3.4 Effect of Silica Application on DTPA Extractable Micronutrients (Fe, Mn, Cu and Zn) in Post-Harvest Soil

The experimental data presented in Table 4 indicates effect of Silica on DTPA extractable micronutrients in post-harvest soils like Fe, Mn, Cu and Zn. DTPA extractable micronutrient analysis indicated that high Fe (78.8ppm) and Mn (1.32ppm) content and low Zn (0.35ppm) and Cu (0.21ppm) content was found in initial soil sample. When the initial analysis value was compared with post-harvest analysis value of the same soil, it was observed that there was a decrease (9.6%) and (21.2%) in Fe (71.2ppm) and Mn content (1.04 ppm) respectively. Whereas an increasing trend was observed in Cu(0.46ppm) and Zn (0.56ppm) content in soil due to application of silica. The study revealed that DTPA extractable micronutrients like Fe and Mn in post-harvest soil showed a reverse trend with increased application of silica.

DTPA extractable Fe in post-harvest soil varied from 71.2 ppm to 79.7 ppm i.e., maximum in 100% STD and minimum in 75% STD+400kg SiO₂/ha. With increased fertilizer application, DTPA extractable Fe increased and reached maximum at T₄ (100% STD). In case of application of silica in combination with 50 %/75 % STD, the Fe content showed a reverse trend with increased application of silica. However, there was no significant difference between the treatments. Though DTPA extractable Fe content was reduced but it was not significantly affected by Si treatments. This is in contradiction with findings of scientists, who reported slight increase in the DTPA-Fe content in soil at harvest of rice plants due to application of silicon [33, 34, 35].

Available Mn in post-harvest soil was recorded minimum (1.04 ppm) in T₁₀ (75% STD +400kg SiO₂/ha) and maximum (1.49 ppm) with T₄ (100% STD). With application of increased fertilizer doses, the available Mn content increased. There was a steady increase in Mn content of soil from Control to 75% STD+400kg SiO₂/ha. Increased application of Silica showed reverse trend in DTPA extractable Mn content ultimately showing a positive impact on reducing Mn toxicity in soil. The decrease in the DTPA-Mn content in soil at harvest of rice plants might be due to efficient utilization of Mn by rice plants. Similar results were also noticed [35].

Table 4. Effect of Silica on Fe, Mn, Cu and Zn in post-harvest soil

Treatment details	DTPA Extractable Micronutrients (ppm)			
	Fe	Mn	Cu	Zn
T ₁ = Absolute Control	74.5	1.30	0.20	0.36
T ₂ = 50% STD	75.3	1.37	0.33	0.42
T ₃ = 75% STD	76.5	1.40	0.34	0.45
T ₄ = 100% STD	79.7	1.49	0.37	0.49
T ₅ = T ₂ + 200kg SiO ₂ /ha	76.6	1.28	0.35	0.48
T ₆ = T ₂ + 300kg SiO ₂ /ha	73.3	1.23	0.38	0.50
T ₇ = T ₂ + 400kg SiO ₂ /ha	72.4	1.05	0.40	0.54
T ₈ = T ₃ + 200kg SiO ₂ /ha	75.9	1.25	0.43	0.48
T ₉ = T ₃ + 300kg SiO ₂ /ha	73.7	1.14	0.43	0.55
T ₁₀ = T ₃ + 400kg SiO ₂ /ha	71.2	1.04	0.46	0.56
SE (M)	-	0.06	0.04	0.05
CD (0.05)	NS	0.17	0.13	0.15

[SE- Standard Error, CD- Critical Difference, NS- Non-Significant]

Table 5. Effect of Silica on grain yield, straw yield, and Harvest Index of paddy

Treatment details	Grain yield(q/ha)	Straw yield (q/ha)	Harvest Index
T ₁ = Absolute Control	28.5	33.0	0.463
T ₂ = 50% STD	32.8	36.8	0.472
T ₃ = 75% STD	33.1	37.7	0.468
T ₄ = 100% STD	36.9	40.8	0.475
T ₅ = T ₂ + 200kg SiO ₂ /ha	33.7	38.0	0.470
T ₆ = T ₂ + 300kg SiO ₂ /ha	35.5	39.1	0.476
T ₇ = T ₂ + 400kg SiO ₂ /ha	34.3	38.4	0.472
T ₈ = T ₃ + 200kg SiO ₂ /ha	34.9	38.8	0.473
T ₉ = T ₃ + 300kg SiO ₂ /ha	36.2	39.6	0.478
T ₁₀ = T ₃ + 400kg SiO ₂ /ha	35.6	39.3	0.476
SE (M)	1.37	1.32	-
CD (0.05)	4.09	3.94	NS

[SE- Standard Error, CD- Critical Difference, NS- Non-Significant]

Available Cu in post-harvest soil varied from 0.20 ppm to 0.46 ppm in which lowest Cu content was observed in T₁ i.e., Absolute control (No Fertilizer) and significantly maximum Cu content (0.46 ppm) was observed in T₁₀(75% STD+400kg SiO₂/ha). With increased fertilizer application the DTPA extractable Cu was increased. All the treatments registered significantly higher extractable Cu over control T₁ except T₂ (50% STD). In case of application of silica in combination with 50% / 75% STD, the Cu content increased with increased application of silica without any significant difference.

Available Zn in post-harvest soil varied from 0.36 ppm to 0.56 ppm so far as different treatments are concerned. DTPA extractable Zn was maximum (0.56 ppm) in T₁₀ (75% STD + 400kg SiO₂/ha) which was almost 56% more than that of control T₁. With increased fertilizer application soil availability of Zn increased. No significant difference was marked in case of application of silica in combination with 50 % / 75 % STD. The study revealed that the DTPA extractable Cu and Zn content increased with increased application of silica. These results are in conformity with the findings [34].

3.5 Effect of Silica on Yield Attributing Characters of Paddy

The experimental data presented in Table 5 and Fig. 1 indicates influence of Silica on plant yield attributing characters like grain yield, straw yield, and Harvest Index. Grain yield, straw yield and harvest index as influenced by application of Silica has been presented in Table 5. There was significant increase in yield over control due to application of different doses of Silica along with 50%/75% STD. Maximum grain and straw yield

i.e., 36.9q/ha and 40.8q/ ha was obtained due to application of 100% STD closely followed by 36.2q/ha and 39.6q/ha due to application of. But Harvest Index was maximum (0.478) in 75%STD+300kg SiO₂/ha. This indicates though 100% STD registered maximum yield but by reducing fertilizer dose by 25% along with supplemental dose of Silica i.e., 300kgSiO₂/ha we could achieve almost equal yield.

Grain yield of rice presented in Table 5 revealed that there was a variation in yield from 28.5q/ha to 36.9q/ha so far as different treatments are concerned. Grain yield was maximum (36.9q/ha) in T₄(100% STD) which was almost 30% more than that of control (28.5q/ha) but it was at par (36.2q/ha) with T₉(75% STD+300kg SiO₂/ha). All the treatments registered significantly higher yield over Control. There was a continuous increase in yield as the dose of the fertilizer increased. Application of silica along with 50% STD/75% STD did not have any significant effect on yield increase however, yield was decreased when silica application went beyond 300 kg/ha. This improvement in grain and straw yields might be due to an enhanced growth, yield components and nutrient uptake of rice with the addition of Si. It was also reported a significant increase in grain and straw yields of rice with increasing Si level [36,15].

Variation in straw yield ranged from 33q/ha to 40.8q/ha. Maximum straw yield was obtained in T₄(100% STD) and minimum in control(33q/ha). Around 24% yield increase was observed due to application of fertilizer i.e., 100% STD over no nutrition. Of course, straw yield in 100% STD (T₄) was at par with 75% STD+300kg SiO₂/ha (T₉). An increased trend was marked when dose of fertilizer was increased. Like grain yield,

application of silica in different dose along with 50%/75% STD did not have any significant effect on yield increase but straw yield was decreased over and above 300 kg SiO₂/ha. The increases in both grain and straw yields might be attributed to the positive effect of Si in increasing growth and yield characteristics [37,15], enhancing the pollen viability and photosynthetic activity [38], reducing abiotic and biotic stress, improving structural support and biomass [39] and improving nutrient uptake [15,40].

Harvest index was almost equal irrespective of treatments imposed. But maximum HI was obtained in T₉ (0.478) which was at par with T₄ (0.475) and T₁₀ (0.476) and minimum in Control (0.463). There was no significant difference

between the treatments so far as HI was concerned. Though Harvest index was maximum due to application of 75% STD+300kg SiO₂/ha but no significant difference was obtained among the treatments. This contradicts the findings of scientists, who reported an increased harvest index of rice with Si application [38,15].

Based on the quadratic regression equation ($y = 0.00020x^2 + 0.05876x + 28.62182$), the grain yield when compared with fertilizer dose (without silica application), it was observed that maxima will not be reached since a is >0 (Fig. 2). However, there was steady increase in grain yield from control treatment to 100% STD and reached maximum i.e., 36.9q/ha.




Fig. 1. Grain yield, straw yield and harvest index




Fig. 2. Maximum yield vs optimum level of fertilizer applied




Fig. 3. Maximum yield vs optimum level Silica applied along with 50% STD fertilizer

Based on the quadratic regression equation ($y = -0.00015x^2 + 0.09300x + 21.1$) the vertex is the highest point on the graph of quadratic function (Fig. 3). The value of vertex ($x = -b/2a$) was 310 with $Y \text{ max} = a (X \text{ max})^2 + b (X \text{ max}) + c = 35.51$. Therefore, 310 kg/ha SiO_2 was the optimum level of Si fertilizer that could provide the maximum yield (35.51q/ha) in combination with 50% STD based on the quadratic regression model. From this point, more amount of Si fertilizer application could result in a decrease in the yield.

Based on the quadratic regression equation ($y = -0.00010x^2 + 0.06050x + 26.6$) the vertex is the highest point on the graph of quadratic function (Fig. 4). The value of vertex ($x = -b/2a$) was 336.1 with $Y \text{ max} = a (X \text{ max})^2 + b (X \text{ max}) + c = 36.76$. Therefore, 336.1 kg/ha SiO_2 was the optimum level of Si fertilizer that could provide the maximum yield (36.76q/ha) in combination with 75% STD based on the quadratic regression model. From this point, more amount of Si fertilizer application could result in a decrease in the yield.




Fig. 4. Maximum yield vs optimum level Silica applied along with 75% STD fertilizer

4. CONCLUSION

Silicon nutrition significantly influenced the growth and yield attributing characteristics as well as the nutrient status in soil. Si is known for its role in alleviating the negative stress effects on many plant species. There was significant increase in number of effective tillers per plant, length of the panicle, grain yield and straw yield over control due to application of silica. However, in case of 1000 grain wt. and Harvest Index, there was no significant difference between the treatments. The study showed increase in pH and Organic Carbon (OC) in post-harvest soil due to application of Silica as compared to initial soil test value. Hence it was suggested to go for application of silica for correcting soil acidity and improving soil OC content in acidic laterite soils. Application of Silica was also effective in reducing the level of Fe and Mn content in post-harvest soil indicated better response in reducing Fe and Mn toxicity in soil. Available N, P, K and S status in post-harvest soil increased significantly due to application of silica along with fertilizer suggested integration of Silica with fertilizer dose without sole application. A reverse trend in DTPA extractable Fe and Mn toxicity was observed in silica application with fertilizer. But DTPA extractable Cu and Zn shows increasing trend due to application of silica. Application of silica along with STD would help in the sustainable production of rice. Future research is needed to find out the heavy metal analysis of the BOF slag to assess its impact on soil due to heavy dose of application. Multilocal trials under different agro-ecosystems with several test crops should be conducted to assess the best possible outcome.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Cuong TX, Ullah H, Datta A, Hanah TC. Effects of Silicon-Based Fertilizer on Growth, Yield and Nutrient Uptake of Rice in Tropical Zone of Vietnam. *Rice Science*, 2017;24(5):283-290.
2. Nagula S, Joseph B and Gladis R. Silicon Nutrition to Rice (*Oryza sativa* L.) Alleviates Fe, Mn and Al Toxicity in Laterite Derived Rice Soils. *Journal of the Indian Society of Soil Science*. 2016;64(3):297-301.
3. Epstein E. Silicon: Its manifold roles in plants. *Ann Appl Biol*. 2009;155(2):155-160.
4. Keeping MG, Reynolds OL. Silicon in agriculture: New insights, new significance and growing application. *Ann Appl Biol*. 2009;155:153-154.
5. Meena VD, Dotaniya ML, Coumar V, Rajendiran S, Kundu S and Rao AS. A case for silicon fertilization to improve crop yields in tropical soils. *Proc Natl. Acad. Sci Ind, Sect B: Biol Sci*. 2014;84:505-518.
6. Farooq MA, Dietz KJ. Silicon as versatile player in plant and human biology: Overlooked and poorly understood. *Front Plant Sci*. 2015;6:1-14.
7. Prakash NB, Nagraj H, Guruswamy KT, Vishwanatha BN, Narayanaswamy C, Gowda NAJ, Vasuki N, Siddaramappa R. Rice hull ash as a source of silicon and phosphatic fertilizers: effect on growth and yield of rice in coastal Karnataka, India. *IRRN*. 2007;32(1):34-36.
8. Snyder GH, Motinchenkov VV, Datnoff LE. Silicon. In *Hand Book of Plant Nutrition* (K.V. Berker and J.P. David, Eds.), Taylor and Francis Group, Broken, Sound Parkway, N.W. 2006;551-568.
9. Rao GB, Yadav P PI, Syriac EK. Effect of Silicon on Soil Physico-chemical Properties in Laterite derived Paddy Soils of Kerala, *J Krishi Vigyan*. 2018;6(2):75-77.
10. Sawant AS, Patil VH, Sawant NK. Rice hull ash applied to seedbed reduces dead hearts in transplanted rice. *IRRN*. 1994; 19(4):21-22.
11. Sudhakar PC, Singh JP, Singh K. Effect of silicon sources and fertility levels on transplanted rice. *IRRN*. 2004;29(2):61-63.
12. Singh AK, Singh R, Singh K. Growth, yield and economics of rice (*Oryza sativa*) as influenced by level and time of silicon application. *Indian Journal of Agronomy*. 2005;50(3):190-193.
13. Singh K, Singh R, Singh KK, Singh Y. Effect of silicon carriers and time of application on rice productivity in a rice-wheat cropping sequence. *IRRN*. 2007; 32(1):30-31.
14. Muriithi C, Mugal E, Kihurani AW, Nafuma CJ, Amboga S. Determination of silicon from rice by-products and chemical sources on rice blast management. 12th

- KARI Scientific conference proceedings; 2010.
15. Pati S, Pal B, Badole S, Hazra GC & Mandal B. Effect of Silicon Fertilization on Growth, Yield, and Nutrient Uptake of Rice, Communications in Soil Science and Plant Analysis. 2016;47(3):284-290.
 16. Shashidhar HE, Chandrashekhar N, Narayanaswamy C, Mahendra AC, Prakash NB. Calcium silicate as silicon source and its interaction with Nitrogen in aerobic rice. IVth Silicon in Agriculture Conference Wild Coast Sun, KwaZulu-Natal, South Africa. 2008;93.
 17. Prabhu AS, Barbosa Filho MP, Filippi MC, Datnoff LE, and Snyder GH. Silicon from rice disease control perspective in Brazil. Studies in Plant Science. 2001;8:293-311.
 18. Dallagnol LJ, Rodrigues FA, Mielli MVB, Ma JF. Rice grain resistance to brown spot and yield are increased by silicon. Tropical Plant Pathology. 2014;39(1):56–63.
 19. Sandhya K. Diatomaceous earth as a source of Silicon on growth and yield of Rice (*Oryza sativa*. L) M.Sc. (Ag) thesis, University of Agricultural Sciences, Bengaluru. 2013;164.
 20. Wallace A. Participation of silicon in cation-anion balance as a possible mechanism for aluminium and iron tolerance in some gramineae. Journal of Plant Nutrition. 1993;16:547-553.
 21. Nwite JC, Obalum SE, Igwe CA, Wakatsuki T. Properties and potential of selected ash sources for improving soil conditions and Sawah rice yields in a degraded in land valley in Southeastern Nigeria. World Journal of Agricultural Sciences. 2011;7(3):304-310.
 22. Qiang FU, Hong H, Ming WU, Zheng CU. Silicon-mediated amelioration of Fe²⁺ toxicity in rice (*Oryza sativa* L.) roots. Pedosphere. 2012;22(6):795-802.
 23. Njoku C, Mbah CN, Okonkwo CI. Effect of rice mill wastes application on selected soil physical properties and maize yield on an ultisol in Abakaliki, Southeastern Nigeria. Journal of Soil Science and Environmental Management. 2011;2(11): 375-383.
 24. Devanur V. Silicon-Solution for tomorrow, Concept note; 2015. Available:<http://www.privilifesciences.com/download/silicon-supplement.pdf>. [25 Dec. 2016]
 25. Burbey A, Rizaldi B, Yulizar Z. Response of upland rice to potassium and silicate application on Ultisol. Pemberitaan Penelitian Sukarami. 1988;15:26-31.
 26. Liang Y. Effects of silicon on enzyme activity and sodium, potassium and calcium concentration in barley under salt stress. Plant Soil. 1999;209(2):217-224.
 27. Mali M, Aery NC. Silicon effects on nodule growth, dry matter production, and mineral nutrition of cowpea (*Vigna unguiculata*). J Plant Nutr. Soil Sci. 2008;171:835-40.
 28. Sadanandan AK, Verghese EJ. Role of Silicate in the uptake of nutrients by rice plants in the lateritic soils of Kerala. Agric. Res. J. Kerala. 1969;7:91-96.
 29. Sawant NK, Snyder GH, Datnoff LE. Silicon management and sustainable rice production. Adv. Agron. 1997;58:151-199.
 30. Selva Kumari G, Baskar M, Jayanthi D, Mathan KK. Effect of integration of fly ash with fertilizers and organic manures on nutrient availability, yield and nutrient uptake at rice in Alfisols, Journal of Indian Society Soil Science. 2000;48(1):34-37.
 31. Subramanian S, Gopalswamy A. Effect of moisture, organic matter, phosphate, and silicate on availability of silicon and phosphorus in rice soils. Journal of the Indian Society Soil Science. 1991;39:99–103.
 32. Chen DQ, Cao BB, Wang SW, Liu P, Deng XP, Yin LA, Zhang SQ. Silicon moderated the K deficiency by improving the plant-water status in sorghum. Scientific Reports. 2016;6:22882.
 33. Sikka R, Kansal BD. Effect of fly-ash application on yield and nutrient composition of rice, wheat and on pH and available nutrient status of soils. Bioresource Tech. 1995;51(2-3):199-203.
 34. Das BK, Choudhury BH and Das KN. Effect of integration of fly ash with fertilizers and FYM on nutrient availability, yield and nutrient uptake of rice in Inceptisols of Assam, India. International Journal of Advanced Research and Technology. 2013;2(11):190-208.
 35. Patil AA. Effect of silicon application along with chemical fertilizers on nutrient uptake and nutrient availability for rice plants, International Journal of Chemical Studies 2018;6(1):260-266.
 36. Deren CW, Datnoff LE, Snyder GH and Martin FG. Silicon concentration, disease response and yield components of rice genotypes grown on flooded organic histosols. Crop Science, 1994;34(3):733–737.

37. Prakash NB, Chandrashekar N, Mahendra C, Patil SU, Thippeshappa GN, Laane HM. Effect of foliar spray of soluble silicic acid on growth and yield parameters of wetland rice in hilly and coastal zone soils of Karnataka, South India. *Journal of Plant Nutrition*. 2011;34(12):1883-1893.
38. Detmann KC, Araújo WL, Martins SCV, Sanglard LMVP, Reis JV, Detmann E, Rodrigues FÁ, Nunes-Nesi A, Fernie AR, DaMatta FM. Silicon nutrition increases grain yield, which, in turn, exerts a feed-forward stimulation of photosynthetic rates via enhanced mesophyll conductance and alters primary metabolism in rice, *The New Phytologist*. 2012;196(3):752-762,11.
39. Meharg C, Meharg AA. Silicon, the silver bullet for mitigating biotic and abiotic stress, and improving grain quality, in rice? *Environmental and Experimental Botany*. 2015;120:8-17.
40. Crooks R, Prentice P. Extensive investigation into field-based responses to a silica fertilizer. *Silicon*. 2017;9(2):301–304.

© 2021 Das et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/77740>